CP3-Origins-2016-005 DNRF90 DIAS-2016-5

Flavor and CP symmetries for leptogenesis and $0\nu\beta\beta$ decay

Claudia Hagedorn and Emiliano Molinaro

CP^3-Origins and Danish Institute for Advanced Study, University of Southern Denmark,
Campusvej 55, DK-5230 Odense M, Denmark

E-mail: hagedorn@cp3.sdu.dk, molinaro@cp3.sdu.dk

arXiv:1602.04206v1 [hep-ph] 12 Feb 2016

Abstract

We perform a comprehensive analysis of the phenomenology of leptonic low and high energy CP phases in a scenario with three heavy right-handed neutrinos in which a flavor and a CP symmetry are non-trivially broken. All CP phases as well as lepton mixing angles are determined by the properties of the flavor and CP symmetry and one free real parameter. We focus on the generation of the baryon asymmetry Y_B of the Universe via unflavored thermal leptogenesis and the predictions of m_{ee}, the quantity measurable in neutrinoless double beta decay. We show that the sign of Y_B can be fixed and the allowed parameter range of m_{ee} can be strongly constrained. We argue on general grounds that the CP asymmetries ϵ_i are dominated by the contribution associated with one Majorana phase and that in cases in which only the Dirac phase is non-trivial the sign of Y_B depends on further parameters. In addition, we comment on the case of flavored thermal leptogenesis where in general the knowledge of the CP phases (and light neutrino mass spectrum) is also not sufficient in order to fix the sign of the CP asymmetries. As examples we discuss the series of flavor groups $\Delta(3\,n^2)$ and $\Delta(6\,n^2)$, $n \geq 2$ integer, and several classes of CP transformations.

1 Introduction

The baryon asymmetry of the Universe Y_B has been precisely measured [1]

$$Y_B = \left. \frac{n_B - n_{\overline{B}}}{s} \right|_0 = (8.65 \pm 0.09) \times 10^{-11} \,. \tag{1}$$

The error is given at the $1\,\sigma$ level and the subscript "0" refers to the present epoch. The generation of Y_B requires the fulfillment of the three Sakharov conditions [2]: C and CP violation, departure from thermal equilibrium and baryon number violation. All three of them are met by the mechanism of (unflavored and flavored) thermal leptogenesis [3]. In fact the decay of right-handed (RH) neutrinos N_i (we always assume the existence of three such states and thus $i = 1, 2, 3$) leads to a lepton asymmetry that is partially converted to a baryonic one via sphaleron processes [4]. Departure from thermal equilibrium arises, since the rate of the Yukawa interactions of RH neutrinos is small compared to the Hubble rate, while complex Yukawa couplings Y_D are responsible for C and CP violation. The relevant quantities for computing Y_B are: the yield of RH neutrinos at high temperatures and the sphaleron conversion factor giving together rise to 10^{-3}, the efficiency factors, taking into account washout and decoherence effects and usually of order of $10^{-3} \div 10^{-1}$, and the CP asymmetries $\epsilon_i^{(\alpha)}$, generated in the decays of the $i^{\rm th}$ RH neutrino N_i and to the flavor $\alpha = e, \mu, \tau$ (if flavored leptogenesis is studied). As has been shown in [5], even in a scenario in which light neutrinos acquire masses via the ordinary type 1 seesaw mechanism [6] the CP phases present in the Yukawa couplings are in general unrelated to the CP phases potentially measurable at low energies: the Dirac phase δ in neutrino oscillation experiments [7–11] and a combination of the two Majorana phases α and β in neutrinoless double beta ($0\nu\beta\beta$) decay [12]. However, it is also well-known that low energy CP phases may be crucial for having successful leptogenesis in case flavor effects are relevant [13].

In theories with flavor G_f and CP symmetries all CP phases can be related. Such symmetries are usually introduced in order to explain the mixing pattern(s) observed in the lepton (and quark) sector. Since two of the lepton mixing angles (and thus most of the entries of the Pontecorvo-Maki-Nakagawa-Sakata (PMNS) mixing matrix U_{PMNS}) are large [14]

$$0.270 \leq \sin^2\theta_{12} \leq 0.344 \,,\ 0.382[9] \leq \sin^2\theta_{23} \leq 0.643[4] \ \text{ and } \ 0.0186[8] \leq \sin^2\theta_{13} \leq 0.0250[1] \tag{2}$$

($3\,\sigma$ ranges for light neutrino masses following normal ordering (NO) and in square brackets for inverted ordering (IO), see also [15, 16]), it is natural to assign the three generations of left-handed (LH) lepton doublets l to an irreducible three-dimensional representation **3** of the flavor symmetry. If the setup contains RH neutrinos, it is reasonable to also assign these to such a representation. Indeed, it is convenient to use the same representation **3** as for l in a non-supersymmetric (non-SUSY) context, while in SUSY models ν_i^c are usually put in the complex conjugated representation with respect to l. The most promising choice of G_f turns out to be a discrete, finite, non-abelian group [17, 18] that is broken to different residual symmetries G_e and G_ν in the charged lepton and neutrino sectors, respectively [19]. These symmetries are in general abelian subgroups of G_f with three different elements at least and, in the particular case of three Majorana neutrinos, that we will consider, $G_\nu = Z_2 \times Z_2$. As is known, with this approach only the Dirac phase can be predicted and the analyses in [20] have shown that it always turns out to be trivial, if mixing angles are required to be in good agreement with experimental data.[1] Thus, in order to fix the Majorana phases via symmetries as well as to achieve δ different from 0 and π this approach has to be modified. An extension that has been proven very powerful [21, 23–28] is to amend the flavor with a CP symmetry [21] (see also [22]). The latter acts in general also in a non-trivial way on the flavor space [29], requiring certain conditions to be fulfilled for having a consistent theory [21, 30, 31]. The choice of G_e remains the same, whereas G_ν is taken to be the direct product of a Z_2 symmetry,

[1] Considering smaller residual symmetries G_e or G_ν or taking into account corrections obviously changes this conclusion, but also reduces the predictive power of the approach.

contained in G_f, and the CP symmetry: $G_\nu = Z_2 \times CP$ [21]. The main feature of such a setting is predicting the mixing angles as well as all three CP phases in terms of only one free real parameter θ that can be chosen without loss of generality to lie in the interval $0 \leq \theta < \pi$. This parameter is present and not fixed in this approach, simply because the residual flavor symmetry in the neutrino sector is only a Z_2 group. On the other hand, involving CP leads to the clear advantage to determine all three CP phases and not only the Dirac phase. Thus, such a scenario is in particular suitable for studying the phenomenology related to CP phases, like leptogenesis and $0\nu\beta\beta$ decay.

Thermal leptogenesis, unflavored as well as flavored, has already been studied in scenarios with the flavor symmetries A_4 and S_4 only [32–38]. One striking feature of these scenarios is the fact that CP asymmetries vanish in case one only considers the leading order (LO) terms in the theory, e.g. those which preserve $G_e = Z_3$ and $G_\nu = Z_2 \times Z_2$ in the charged lepton and neutrino sectors, respectively. Thus, taking into account next-to-leading order (NLO) corrections (in the neutrino sector, in particular, to Y_D) is mandatory. These are in general proportional to the (small) symmetry breaking parameter $\kappa \sim 10^{-(2\div 3)}$. As shown in [33], the CP asymmetries ϵ_i are in general proportional to κ^2 for unflavored leptogenesis, whereas in the case of flavored leptogenesis the suppression is smaller and ϵ_i^α are proportional to κ only.

$0\nu\beta\beta$ decay has already been discussed in scenarios with a flavor and a CP symmetry, see last reference in [23], first reference in [28] as well as first reference in [24], [39] and [26]. In [39] the authors have considered the series of groups $\Delta(6\,n^2)$, but they have assumed that the residual symmetry in the neutrino sector is larger $G_\nu = Z_2 \times Z_2 \times CP$. Thus, the Dirac phase is fixed to be trivial as well as one of the Majorana phases which clearly affects the predictions for m_{ee}. In [26] the authors instead study $G_\nu = Z_2 \times CP$ as residual group like in our analysis. However, the presented results are mostly in the limit in which the group theoretical parameter n is taken to be very large and no particular choice of the residual CP symmetry is made.

In the present paper we study leptogenesis and $0\nu\beta\beta$ decay in a scenario with the flavor symmetry $\Delta(3\,n^2)$ or $\Delta(6\,n^2)$ and a CP symmetry that are broken in a non-trivial way. As discussed in [26,27], for several choices of flavor groups $\Delta(3\,n^2)$ or $\Delta(6\,n^2)$, CP symmetries, residual groups G_e and G_ν and the free parameter θ it is possible to obtain lepton mixing angles in agreement with the experimental data and in turn to predict the CP phases δ and α, β. We base our study on these results and we assume throughout that the charged lepton mass matrix is governed by the residual symmetry $G_e = Z_3$, while the RH neutrino mass matrix is taken to be of the most general form compatible with $G_\nu = Z_2 \times CP$. Since LH lepton doublets and RH neutrinos transform according to the same three-dimensional representation **3** of the flavor group in a non-SUSY framework (or in a SUSY context l and ν^c as $\overline{\mathbf{3}}$ and **3**), the Yukawa coupling Y_D of neutrinos with trivial flavor structure is invariant under the entire flavor group and CP. We note that RH neutrinos are not strongly hierarchical in our scenario and hence all three of them are expected to be relevant for leptogenesis. We mainly focus on the case of unflavored leptogenesis and only highlight the main differences occurring, if the baryon asymmetry of the Universe is instead generated via flavored leptogenesis. Similarly, most of our results for leptogenesis are obtained in a non-SUSY framework, however, we emphasize the changes that have to be implemented in order to apply these to a SUSY model. We find that the CP asymmetries ϵ_i vanish for unflavored leptogenesis, if we only consider LO terms. Thus, we have to rely on NLO terms. In a generic model these can be arbitrary, but frequently it turns out that one type is dominating and that it is still invariant under one of the residual symmetries. Here we consider a case in which the dominant NLO contribution arises in the neutrino sector and only corrects Y_D. Furthermore, we assume that this correction δY_D stems from the charged lepton sector and is thus constrained by the residual group $G_e = Z_3$. As expansion parameter we use κ and hence $\delta Y_D \propto \kappa$. It induces an appropriate suppression of the CP asymmetries $\epsilon_i \propto \kappa^2$, similar to what has been already observed in scenarios with a flavor symmetry only. We show that the sign of ϵ_i is fixed depending on the low energy CP phases. In particular, the phases, contained in δY_D, are irrelevant for the LO result of ϵ_i. The sign of Y_B is determined as well and it is given by ϵ_i and the sign of the loop function, traced back to the light neutrino mass spectrum that can be hierarchical with NO, IO or quasi-degenerate (QD). We do not only perform a comprehensive analytical study in which we consider the different choices of residual symmetries,

found in [27], but we also consider the case in which the mixing in the RH neutrino sector is given by a general form of the PMNS mixing matrix. The latter study reveals three interesting features. The CP asymmetries ϵ_i can be expressed in terms of the CP invariants I_i that are proportional to $\sin\alpha$, $\sin\beta$ and $\sin(\alpha-\beta)$ (for definition see appendix A.1). The dominant contribution is expected to arise from terms proportional to $\sin\alpha$, as long as this Majorana phase does not take a value close to 0 or π or the loop function suppresses these terms for particular values of the light neutrino masses. In case $\sin\delta$ is dominating the CP asymmetries ϵ_i their sign cannot be predicted and depends on the relative size of the parameters appearing in δY_D as well as on the octant of the atmospheric mixing angle θ_{23}. We exemplify these findings with several cases and detail instances in which the sign of Y_B is correctly determined. Furthermore, we comment on the case of flavored leptogenesis. The CP asymmetries ϵ_i^α vanish as well, if we only take into account the LO terms, since the Yukawa couplings of neutrinos are taken to be invariant under the flavor symmetry G_f and CP in our analysis. Also in this case corrections $\delta Y_D \propto \kappa$ induce non-zero ϵ_i^α. However, their sign depends in general on the parameters contained in δY_D. A way to generate $\epsilon_i^\alpha \neq 0$ without corrections δY_D is to assume the Yukawa couplings of neutrinos to be of the most general form compatible with the residual symmetry $G_\nu = Z_2 \times CP$. Again, the sign of ϵ_i^α depends on parameters that are not directly related to the low energy CP phases (and the light neutrino mass spectrum). So, we arrive at the conclusion that the sign of the CP asymmetries cannot be univocally predicted in the flavored regime. Regarding $0\nu\beta\beta$ decay we carefully study it analytically for different choices of symmetries G_f, G_e and G_ν and present several numerical examples, pointing out the following interesting features: for neutrino masses following IO it is possible to achieve only values close to the very upper limit $m_{ee} \approx 0.05\,\text{eV}$ of the parameter space, generally compatible with experimental data, and thus enhancing the prospects for a discovery of $0\nu\beta\beta$ decay in the not-too-far future; for normally ordered light neutrino masses we observe that the well-known cancellation of the different terms contributing to m_{ee} can be avoided and thus a lower limit on $m_{ee} \gtrsim 10^{-3}\,\text{eV}$ can be set or at least the interval of the lightest neutrino mass m_0 for which a noticeable cancellation in m_{ee} occurs to be considerably shrunk.

The paper is structured as follows: in section 2 we describe our scenario in which a flavor group $G_f = \Delta(3\,n^2)$ and $G_f = \Delta(6\,n^2)$ combined with a CP symmetry is broken to $G_e = Z_3$ and $G_\nu = Z_2 \times CP$ and corrections from the charged lepton sector to the neutrino one are crucial for the generation of non-vanishing CP asymmetries. We also briefly repeat the results obtained in [27] for lepton mixing, in particular for the CP invariants, in section 2. Section 3 contains our analysis of unflavored thermal leptogenesis in our scenario, first mentioning some general properties of the underlying leptogenesis framework, then discussing the case in which the mixing matrix U_R, diagonalizing the RH neutrino mass matrix, is taken to be of the general form of the PMNS mixing matrix, and afterwards analytically the properties of the scenarios in which U_R is determined by the breaking of G_f and the CP symmetry (distinguishing the different cases case 1) through case 3 b.1) and case 3 a) that give rise to different results for lepton mixing) and eventually also numerically. We emphasize the cases in which the sign of the CP asymmetries (and thus of the baryon asymmetry of the Universe) that is intimately related to the CP phases δ, α and β can be predicted and also point out in which cases this is in general impossible. In section 4 we discuss flavored thermal leptogenesis and show with several examples that the sign of the CP asymmetries ϵ_i^α can in general not be predicted just from the knowledge of the low energy CP phases and the light neutrino mass spectrum. We also demonstrate that for Y_D invariant under G_ν (and not G_f and CP) non-zero CP asymmetries can be achieved without corrections δY_D. We comment on the changes occurring, if our scenario is realized in a SUSY context in section 5 and point out the similarities and crucial differences. Section 6 is dedicated to an analytical and numerical study of $0\nu\beta\beta$ decay in which the effects of constraining the lepton mixing parameters, in particular the two Majorana phases α and β, becomes apparent. The summary of our results can be found in section 7. Our conventions for mixing angles θ_{ij} and the CP phases δ, α and β together with the results of the global fit [14] are given in appendix A. The choice of generators of the flavor groups $\Delta(3\,n^2)$ and $\Delta(6\,n^2)$ are presented in appendix B. For completeness, in appendix C some results for ϵ_i are shown for the case in which U_R is identified with the general PMNS mixing matrix. Appendix D contains sketches

of explicit models which realize the envisaged scenario in a non-SUSY as well as a SUSY context, choosing $G_f = \Delta(6\,n^2)$ with $n = 4$ and for the parameter s characterizing the CP symmetry $s = 1$.

2 Approach and lepton mixing resulting from $G_f = \Delta(3\,n^2)$ and $G_f = \Delta(6\,n^2)$ and CP

Here we present the approach we follow and the results obtained for lepton mixing from the groups $G_f = \Delta(3\,n^2)$ and $\Delta(6\,n^2)$ discussed in [27].

2.1 Approach

We consider in the following a scenario with three RH Majorana neutrinos N_i. These form a faithful and irreducible representation **3** of the flavor group G_f. To the very same three-dimensional representation we also assign the three generations of LH leptons l.[2] The RH charged leptons α_R, $\alpha = e,\ \mu,\ \tau$ are all assumed to transform trivially under G_f, i.e. $\alpha_R \sim \mathbf{1}$ under G_f. In order to distinguish them we employ an auxiliary symmetry $Z_3^{(\mathrm{aux})}$ under which they carry the charges 1, ω and ω^2. The phase ω is defined as the third root of unity: $\omega = e^{2\,\pi\,i/3}$. LH leptons and RH neutrinos are instead neutral under $Z_3^{(\mathrm{aux})}$. In addition, the theory is invariant under a CP symmetry whose action is represented by the CP transformation $X_{\mathbf{r}}$ in the different representations $\mathbf{r}$ of G_f. In general, this CP symmetry acts also non-trivially on the flavor space [29] (see also [30]). The matrix $X_{\mathbf{r}}$ is unitary and symmetric (for details see [21])

$$X_{\mathbf{r}} X_{\mathbf{r}}^{\dagger} = X_{\mathbf{r}} X_{\mathbf{r}}^{\star} = 1\ . \tag{3}$$

As shown in [21], the consistent combination of a flavor group G_f and a CP symmetry, represented by $X_{\mathbf{r}}$, requires that the condition

$$(X_{\mathbf{r}}^{-1} g_{\mathbf{r}} X_{\mathbf{r}})^{\star} = g'_{\mathbf{r}} \tag{4}$$

is fulfilled for all elements g of G_f with g' being also an element of G_f that is in general different from g. Here $g_{\mathbf{r}}$ and $g'_{\mathbf{r}}$ indicate that both elements, g and g', are given in the representation $\mathbf{r}$. Since we only make explicit use of the trivial representation and the representation **3** (and its complex conjugate), we only need to specify the form of the CP transformation in these two representations. Without loss of generality we can choose $X_{\mathbf{1}} = 1$, while the form of $X \equiv X_{\mathbf{3}}$ is in general non-trivial. Its particular form is given explicitly in the different cases, see (28). Similarly, we only need to consider the representation matrices $g_{\mathbf{r}}$ of the abstract elements g of the group G_f in the trivial representation and in **3**. Those belonging to the former representation always equal the identity, $g_{\mathbf{1}} = 1$, while we denote those of the latter, for simplicity, with the same symbol as the abstract elements themselves, i.e. $g_{\mathbf{3}} \equiv g$.[3]

We focus on a non-SUSY framework (and comment on the implementation in a SUSY context in section 5 and appendix D) in the present paper and thus the form of the relevant Lagrangian is

$$\mathcal{L}_l = -Y_l\,\bar{l}\,H\,\alpha_R - Y_D\,\bar{l}\,H^c N - \frac{1}{2}\overline{N^c} M_R N + \mathrm{h.c.} \tag{5}$$

with H being the Higgs $SU(2)_L$ doublet and $H^c = \epsilon\,H^{\star}$. Note that the Higgs field does neither transform under the flavor nor the CP symmetry of the theory.

The residual symmetry in the charged lepton sector is chosen as

$$G_e = Z_3^{(D)}\ , \tag{6}$$

[2] In the SUSY model we choose l and ν^c to transform in two complex conjugated three-dimensional representations of the group G_f, see section 5.

[3] In appendix D we discuss a sketch of a model. This requires the knowledge of the representation matrices in further representations of the flavor group as well as of the form of the CP transformation in the other representations. However, these pieces of information are either available in the literature or can be straightforwardly calculated.

where $Z_3^{(D)}$ is the diagonal subgroup of the Z_3 symmetry contained in G_f and the additional Z_3 group $Z_3^{(\text{aux})}$. The requirement to preserve G_e in the charged lepton sector is equivalent to requiring that the Yukawa matrix Y_l is invariant under this symmetry. For convenience, we choose a basis for all groups G_f we discuss in such a way that the generator Q of the subgroup Z_3 contained in G_f is diagonal. Thus, the charged lepton mass matrix m_l is diagonal and it contains three independent parameters that can be identified with the charged lepton masses

$$m_l = Y_l \, v_{\text{EW}} = \text{diag}\,(m_e, m_\mu, m_\tau) \; . \tag{7}$$

The value of the Higgs VEV is fixed to $v_{\text{EW}} = \langle H \rangle \approx 174\,\text{GeV}$ in our convention. If the ordering of m_e, m_μ and m_τ is not the standard one, we apply a permutation matrix P. Thus, the contribution of the charged leptons to the PMNS mixing matrix is in the case at hand given by $U_l = P$ (up to unphysical phases). In an explicit model the matrix m_l can be generated when appropriate flavor symmetry breaking fields take a vacuum expectation value (VEV) leaving invariant $Z_3^{(D)}$, see appendix D. In such a realization also the mass hierarchy of the charged leptons is usually explained either with the help of an additional Froggatt-Nielsen symmetry $U(1)_{\text{FN}}$ [40] under which RH charged leptons carry different charges, see e.g. [41], or it is generated through operators with multiple insertions of flavor symmetry breaking fields, see e.g. [42]. Notice the additional symmetry $Z_3^{(\text{aux})}$, not present in the model-independent approach [27], does not have a direct impact on our results on lepton mixing and thus the results in subsection 2.2 hold without loss of generality.

Since LH leptons and RH neutrinos transform in the same way under all flavor and CP symmetries, the second term in (5) is automatically invariant under these symmetries and the Yukawa coupling Y_D is proportional to the identity matrix in flavor space. The Dirac mass matrix of the neutrinos hence reads

$$m_D = Y_D \, v_{\text{EW}} = y_0 \, v_{\text{EW}} \, \mathbb{1} \; . \tag{8}$$

It is clear that the form of m_D preserves G_f and the phase of the parameter y_0 can always be chosen in such a way that the imposed CP symmetry remains intact. Throughout this paper we take y_0 to be positive, since the phase of y_0 can be absorbed in a re-definition of fields and is thus unphysical. Clearly, m_D is, as consequence, also invariant under all subgroups of G_f and CP.

The presence of the non-trivial residual symmetry

$$G_\nu = Z_2 \times CP \tag{9}$$

that is the direct product of a Z_2 subgroup of the flavor symmetry G_f and the CP symmetry instead manifests itself in the form of the Majorana mass matrix M_R of RH neutrinos. Before presenting its explicit form we mention that the generator of the Z_2 group, denoted by Z in **3**, and the CP transformation X, have to fulfil[4]

$$X Z^\star - Z X = 0 \; , \tag{10}$$

since the product of these two symmetries is required to be a direct one. Furthermore, we note that also ZX is a CP symmetry present in the neutrino sector, satisfying the condition in (10) [21]. The Majorana mass matrix M_R of RH neutrinos is assumed to be invariant under the residual group G_ν, i.e.

$$U_R^T \, M_R \, U_R = \text{diag}\,(M_1, M_2, M_3) \tag{11}$$

with M_i being the three masses and the unitary matrix U_R is of the form

$$U_R = \Omega \, R_{ij}(\theta) \, K_\nu \; . \tag{12}$$

The matrix Ω is determined by X and Z, $R_{ij}(\theta)$ is a rotation matrix in the (ij)-plane through the real parameter θ, $0 \leq \theta < \pi$, and K_ν is a diagonal matrix with elements ± 1 and $\pm i$. The latter is necessary for making the masses of RH neutrinos real and positive and we parametrize it as

$$K_\nu = \begin{pmatrix} 1 & 0 & 0 \\ 0 & i^{k_1} & 0 \\ 0 & 0 & i^{k_2} \end{pmatrix} \quad \text{with} \quad k_{1,2} = 0, 1, 2, 3 \; . \tag{13}$$

[4]This equation is trivially fulfilled, if we consider the trivial representation of G_f instead of **3**.

Thus, the matrix M_R contains in general four independent real parameters: the RH neutrino masses M_i and θ. In an explicit model M_R is generated, like the charged lepton mass matrix m_l, via the couplings of RH neutrinos to flavor symmetry breaking fields. Obviously, the VEVs of the latter should leave the residual symmetry G_ν invariant. The light neutrino masses originate from the type 1 seesaw mechanism [6]

$$m_\nu = -m_D\, M_R^{-1}\, m_D^T = -y_0^2\, v^2\, M_R^{-1}\ . \tag{14}$$

This matrix is diagonalized by U_ν

$$U_\nu^\dagger\, m_\nu\, U_\nu^\star = \text{diag}\,(m_1, m_2, m_3) \quad \text{with} \quad U_\nu = U_R\ . \tag{15}$$

The light neutrino masses m_i are inversely proportional to the RH neutrino masses M_i

$$m_i = -\frac{y_0^2\, v^2}{M_i}\ . \tag{16}$$

Note that we will omit the minus sign appearing in (16) in the following. The matrix $U_\nu^\star$ appears in (15), since the light neutrino mass matrix is given in the basis $\overline{\nu_L}\, m_\nu\, \nu_L^c$ and we identify U_ν with the matrix that brings the fields ν_L into their mass basis so that the lepton mixing matrix U_{PMNS} is given by

$$U_{PMNS} = U_l^\dagger\, U_\nu = P^T U_R = P^T \Omega\, R_{ij}(\theta)\, K_\nu\ . \tag{17}$$

In accordance to what has been shown in [21], in such an approach the form of the PMNS mixing matrix is fixed by the symmetries G_f, G_e and G_ν up to the free real parameter θ, possible permutations of rows and columns, since the masses of charged leptons and neutrinos are not predicted, and possible, but unphysical, phases. Consequently, all mixing angles and CP phases are strongly correlated, because they all only depend on θ. The latter has to be fixed to a particular value in order to accommodate the data on lepton mixing angles well that has to be explained by some mechanism in an explicit model, see e.g. [23].

As mentioned, masses are generally not predicted in this approach, unless a particular model is considered, see e.g. second reference in [23]. Thus, we parametrize the three light neutrino masses in terms of the two measured mass squared differences

$$\Delta m^2_{\text{sol}} = m_2^2 - m_1^2 \quad \text{and} \quad \Delta m^2_{\text{atm}} = \begin{cases} m_3^2 - m_1^2 & \text{for normal ordering (NO)} \\ m_3^2 - m_2^2 & \text{for inverted ordering (IO)} \end{cases} \tag{18}$$

and the lightest neutrino mass m_0. For NO the spectrum is thus of the form

$$m_1 = m_0\ \ , \quad m_2 = \sqrt{m_0^2 + \Delta m^2_{\text{sol}}}\ \ , \quad m_3 = \sqrt{m_0^2 + \Delta m^2_{\text{atm}}}\,, \tag{19}$$

while we find for IO

$$m_1 = \sqrt{m_0^2 + |\Delta m^2_{\text{atm}}| - \Delta m^2_{\text{sol}}}\,, \quad m_2 = \sqrt{m_0^2 + |\Delta m^2_{\text{atm}}|}\,, \quad m_3 = m_0\ . \tag{20}$$

If $m_0 \gtrsim 0.1\,\text{eV} > \sqrt{\Delta m^2_{\text{atm}}}$, the spectrum is called quasi-degenerate (QD) and $m_1 \approx m_2 \approx m_3$. The current best fit values obtained from the global fit analysis in [14] are

$$\begin{aligned} \Delta m^2_{\text{sol}} &= 7.50 \times 10^{-5}\ \text{eV}^2\ , \\ \Delta m^2_{\text{atm}} = 2.457 \times 10^{-3}\ \text{eV}^2\ \ (\text{NO}) \quad &\text{and} \quad \Delta m^2_{\text{atm}} = -2.449 \times 10^{-3}\ \text{eV}^2\ \ (\text{IO})\ . \end{aligned} \tag{21}$$

For the $3\,\sigma$ intervals of the mass squared differences see appendix A.2. The sum of the three light neutrino masses is constrained by cosmology and an upper bound is given by the Planck collaboration (using TT, TE, EE and lowP constraints) [1]

$$\sum_{k=1}^{3} m_k < 0.492\ \text{eV} \quad \text{at 95\% C.L.} \tag{22}$$

As a consequence the lightest neutrino mass m_0 has to be smaller than

$$m_0 \lesssim 0.164 \text{ eV} . \tag{23}$$

Using (16) we see that the masses of the RH neutrinos are determined by the light neutrino mass ordering and by the lightest neutrino mass m_0 and thus fixed once the former are fixed.

The leading order (LO) results are in general perturbed in an explicit model, e.g. the Dirac mass matrix of the neutrinos can receive contributions from flavor symmetry breaking fields dominantly coupling to charged leptons. Thus, we will have in general

$$\begin{aligned} Y_l &= Y_l^0 + \delta Y_l + \dots , \\ Y_D &= Y_D^0 + \delta Y_D + \dots , \quad M_R = M_R^0 + \delta M_R + \dots \end{aligned} \tag{24}$$

with Y_l^0, Y_D^0, M_R^0 denoting the LO results, given in (7), (8) and (11). The corrections δY_l, δY_D and δM_R are suppressed with respect to the LO results by (powers of) the small (real, positive) parameter κ, $\kappa \ll 1$. For simplicity, we include this suppression factor in the definition of the corrections. The latter are responsible for changes in the matrices that diagonalize m_l, M_R and m_ν, i.e. the mixing matrices will read

$$U_l = P\,\delta U_l , \;\; U_R = U_R^0\,\delta U_R \;\; \text{and} \;\; U_\nu = U_\nu^0\,\delta U_\nu \;\; \text{with} \;\; \delta U_l \approx \mathbb{1} , \;\; \delta U_R \approx \mathbb{1} , \;\; \delta U_\nu \approx \mathbb{1} \tag{25}$$

leading to a PMNS mixing matrix of the form

$$U_{PMNS} = U_l^\dagger\, U_\nu = (\delta U_l)^\dagger\, P^T U_\nu^0\, \delta U_\nu \approx P^T U_\nu^0 = U_{PMNS}^0 . \tag{26}$$

In the discussion of leptogenesis in our scenario we are particularly interested in the corrections δY_D to the Yukawa coupling of the neutrinos. There are three principle possibilities: either the dominant correction δY_D arises from fields coupling dominantly to RH neutrinos, then δY_D is also invariant under G_ν, or δY_D arises from fields coupling dominantly to charged leptons, then δY_D is expected to possess as residual symmetry $G_e = Z_3^{(D)}$ or δY_D respects none of these residual symmetries, since combinations of both types of flavor symmetry breaking fields contribute to the dominant correction. A particularly interesting case is the second one. We can parametrize the correction δY_D in this case as

$$\delta Y_D = \begin{pmatrix} \frac{2}{\sqrt{3}} z_1 & 0 & 0 \\ 0 & -\frac{1}{\sqrt{3}} z_1 - z_2 & 0 \\ 0 & 0 & -\frac{1}{\sqrt{3}} z_1 + z_2 \end{pmatrix} \kappa \tag{27}$$

with $z_{1,2}$ being complex numbers with absolute values of order one. These parameters are in general complex, because we do not require a residual CP symmetry to be preserved in the charged lepton sector. Note that the trace of δY_D vanishes, since the correction proportional to the trace can be absorbed in the LO term. Note further that the corrections are normalized in such a way that the trace of the square of the matrices that accompany z_1 and z_2, respectively, is the same, while the trace of the product of these two matrices vanishes. In section 3 we focus on leptogenesis in a scenario with δY_D as in (27).

We present a sketch of a non-SUSY model in appendix D, in particular, in order to motivate the size of the parameter κ that we will assume later in our phenomenological study, see (79).

2.2 Lepton mixing from $G_f = \Delta(3\,n^2)$ and $G_f = \Delta(6\,n^2)$ and CP

In [27] the mixing patterns that can – for certain values of the parameters of the theory – fit the experimental data on the lepton mixing angles well [14] have been found in a scenario with $G_f = \Delta(3\,n^2)$ or $G_f = \Delta(6\,n^2)$, n being an integer, and CP. Like in [27], we will require in the following that three does not divide n as well as for case 1) and case 2) also that n is even. The residual symmetries in the charged lepton and neutrino sectors are $G_e = Z_3$ and $G_\nu = Z_2 \times CP$. As regards the mixing there is no difference in considering as residual symmetry $G_e = Z_3$ generated

by $Q = a$ (see definition of a in appendix B) or the residual symmetry $G_e = Z_3^{(D)}$ where $Z_3^{(D)}$ is the diagonal subgroup of the Z_3 symmetry generated by $Q = a$ and the additional Z_3 group, since the relevant property, namely the fact that the charged lepton mass matrix is diagonal, is not altered. In [27] all possible choices of Z_3 and Z_2 groups and a certain set of CP transformations X have been studied.[5] In particular, the following representative cases that lead to different mixing patterns have been singled out

$$\begin{array}{llll} \Delta(3\,n^2)\,,\ \Delta(6\,n^2) & \text{case 1)} & Z = c^{n/2} & X = a\,b\,c^s\,d^{2s}\,P_{23}\,, \\ \Delta(3\,n^2)\,,\ \Delta(6\,n^2) & \text{case 2)} & Z = c^{n/2} & X = c^s\,d^t\,P_{23}\,, \\ \Delta(6\,n^2) & \text{case 3 a) and case 3 b.1)} & Z = b\,c^m\,d^m & X = b\,c^s\,d^{n-s}\,P_{23}\,, \end{array} \tag{28}$$

with s, t, m taking integer values in the interval $0 \leq s, t, m \leq n-1$, a, c and d (and b) being the generators of the group $\Delta(3\,n^2)$ (and $\Delta(6\,n^2)$), see appendix B and [27], and the matrix P_{23} reading

$$P_{23} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}\,. \tag{29}$$

We note that the CP transformation $X = P_{23}$ in the representation **3** corresponds to the following automorphism acting on the generators a, b, c and d as

$$a \ \to \ a\ ,\ b \ \to \ b\ ,\ c \ \to \ c^{-1} \ \text{ and } \ d \ \to \ d^{-1}\,. \tag{30}$$

Since in our scenario only the Majorana mass matrix of the RH neutrinos is non-diagonal, mixing solely originates from this sector. As shown in the preceding subsection, the diagonalization matrix U_R and thus also the lepton mixing matrix U_{PMNS} are given at LO by the matrix in (12), up to permutations of rows and columns.

The case $X = P_{23}$ in the charged lepton mass basis is well-known in the literature [22] and is called $\mu\tau$ reflection symmetry. Its predictions are maximal atmospheric mixing and maximal Dirac phase as well as trivial Majorana phases, while the solar mixing angle remains in general unconstrained without additional symmetries.

In the following we repeat the form of the CP invariants J_{CP}, I_1 and I_2 for the cases in (28), as these are relevant for our analysis of leptogenesis. We note that we add the results of a third Majorana CP invariant, called I_3 and defined in (171) in appendix A.1. Clearly, this third invariant is not independent of the two other ones I_1 and I_2. However, it proves useful to employ I_3 in the discussion of leptogenesis, see e.g. formula in (75-77). For the formulae of the mixing angles we refer the reader to [27].

In case 1) the PMNS mixing matrix is of the form

$$U_{PMNS,1} = U_{R,1} = \Omega_1\,R_{13}(\theta)\,K_\nu\,, \tag{31}$$

where

$$\Omega_1 = e^{i\,\phi_s}\,U_{TB}\,\begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{-3\,i\,\phi_s} & 0 \\ 0 & 0 & -1 \end{pmatrix} \quad \text{with} \quad \phi_s = \frac{\pi\,s}{n}\,, \tag{32}$$

$$U_{TB} = \begin{pmatrix} \sqrt{\frac{2}{3}} & \frac{1}{\sqrt{3}} & 0 \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{6}} & \frac{1}{\sqrt{3}} & -\frac{1}{\sqrt{2}} \end{pmatrix} \quad \text{and} \quad R_{13}(\theta) = \begin{pmatrix} \cos\theta & 0 & \sin\theta \\ 0 & 1 & 0 \\ -\sin\theta & 0 & \cos\theta \end{pmatrix}\,. \tag{33}$$

The results for the CP invariants are

$$\begin{aligned} & J_{CP} = 0\,, \quad I_1 = \frac{2}{9}\,(-1)^{k_1+1}\,\cos^2\theta\,\sin 6\,\phi_s\,, \quad I_2 = 0\,, \\ & \text{and} \quad I_3 = \frac{2}{9}\,(-1)^{k_1+k_2}\,\sin^2\theta\,\sin 6\,\phi_s\,. \end{aligned} \tag{34}$$

[5] These results have been confirmed in [26] and there also extended to the case in which G_e is not a Z_3 symmetry.

In this case also the form of the Majorana phase α is particularly simple

$$\sin\alpha = (-1)^{k_1+1}\,\sin 6\,\phi_s \tag{35}$$

and we see immediately that for the choices $s = 0$ and $s = n/2$ (i.e. if the CP transformations $X = a\,b\,P_{23}$ and $X = a\,b\,c^{n/2}\,P_{23}$ are imposed, respectively) also the Majorana phase α is trivial, meaning that CP is conserved in this case, see also [27]. We note that replacing θ with $\pi - \theta$ leads to the same results for the solar and reactor mixing angle as well as the in general non-trivial Majorana phase α, whereas the atmospheric mixing angle changes its octant.

In case 2) the matrix U_{PMNS} reads

$$U_{PMNS,2} = U_{R,2} = \Omega_2\,R_{13}(\theta)\,K_\nu \tag{36}$$

with

$$\Omega_2 = e^{i\,\phi_v/6}\,U_{TB}\,R_{13}\left(-\frac{\phi_u}{2}\right)\,\begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{-i\,\phi_v/2} & 0 \\ 0 & 0 & -i \end{pmatrix} \quad \text{with} \quad \phi_u = \frac{\pi u}{n} \quad \text{and} \quad \phi_v = \frac{\pi v}{n}\,. \tag{37}$$

The integer parameters u and v depend on the choice of the CP transformation $X = c^s d^t\,P_{23}$ and are related to the exponents s and t as follows

$$u = 2\,s - t \quad \text{and} \quad v = 3\,t\,. \tag{38}$$

Since $0 \leq s, t \leq n-1$, we get for u and v as admitted intervals: $-(n-1) \leq u \leq 2\,(n-1)$ and $0 \leq v \leq 3\,(n-1)$. According to [27] the CP invariants are of the form

$$\begin{aligned} J_{CP} &= -\frac{\sin 2\theta}{6\sqrt{3}}\;\;,\;\; I_2 = \frac{1}{9}(-1)^{k_2}\,\sin 2\phi_u \sin 2\theta\;\;, \\ I_1 &= \frac{1}{9}(-1)^{k_1+1}\,\left([\cos\phi_u + \cos 2\theta]\sin\phi_v - \sin\phi_u\cos\phi_v\sin 2\theta\right)\;\;, \end{aligned} \tag{39}$$

and the third Majorana CP invariant is given by

$$I_3 = \frac{1}{9}(-1)^{k_1+k_2}\,\left([\cos\phi_u - \cos 2\theta]\sin\phi_v + \sin\phi_u\cos\phi_v\sin 2\theta\right)\;. \tag{40}$$

We can derive approximate formulae for the CP phases [27], if we take into account the constraints on the group theoretical quantities u and n as well as on the free parameter θ arising from the requirement to describe the experimentally measured values of the lepton mixing angles well. In particular, we see that the sine of the Majorana phase α can be expressed as

$$\sin\alpha \approx (-1)^{k_1+1}\,\sin\phi_v\,. \tag{41}$$

The study of several explicit cases in [27] has demonstrated that this formula holds to very good approximation for all values of the parameter v. Thus, we see that choices of $v \approx 0, n, 2\,n$ lead to a very small phase α, while values of $v \approx n/2, 3\,n/2, 5\,n/2$ entail (almost) maximal α, see also table 2 in our numerical discussion in subsection 3.4.

In addition, we note that by permuting the rows of the mixing matrix $U_{PMNS,2} = U_{R,2}$, defined in (36), i.e. we consider either

$$P_1\,\Omega_2\,R_{13}(\theta)\,K_\nu \quad \text{or} \quad P_1^2\,\Omega_2\,R_{13}(\theta)\,K_\nu \tag{42}$$

using the permutation matrix P_1

$$P_1 = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}\;, \tag{43}$$

two slightly different mixing patterns can be obtained whose results for the CP invariants can be deduced from those given in (39-40), if we apply either the transformations

$$u \;\to\; u - \frac{n}{3} \;,\; \theta \;\to\; \frac{\pi}{2} - \theta \;\text{ and }\; k_1 \;\to\; k_1 + 1 \;, \tag{44}$$

or

$$u \;\to\; u + \frac{n}{3} \;,\; \theta \;\to\; \frac{\pi}{2} - \theta \;\text{ and }\; k_1 \;\to\; k_1 + 1 \;. \tag{45}$$

For details see [27]. We only notice here that applying any of the two sets of transformations also changes the sign in the approximate formula for the sine of the Majorana phase α given in (41). In this case replacing θ with $\pi - \theta$ does not affect the mixing angles, whereas the sign of J_{CP} and I_2 change and the CP invariants I_1 and I_3 do not transform in a definite way under this replacement. Nevertheless, using the approximate relation in (41) for $\sin\alpha$, we see that does not change, if we only replace θ with $\pi - \theta$. The two CP invariants I_1 and I_3 can be made transforming in a definite way by additionally sending k_1 into $k_1 + 1$ and v into $k\,n - v$ (with k being an odd integer chosen in such a way that $k\,n - v$ is still in the admitted interval for the parameter v). Then, both, I_1 and I_3, change sign so that all CP invariant change sign. This statement is in agreement with the observation that $\sin\alpha$ in (41) is not affected by the replacement of θ by $\pi - \theta$ alone, but it changes sign, if we change k_1 into $k_1 + 1$.

The third case can be divided into two subcases, case 3 a) and case 3 b.1). Since we show numerical results only for case 3 b.1) in our analysis of leptogenesis, we first present the findings of [27] for this case. For case 3 b.1) the matrix U_{PMNS} takes the form

$$U_{PMNS,3b1} = U_{R,3b1} = \Omega_3\, R_{12}(\theta)\, K_\nu\, P_1 \tag{46}$$

with

$$\Omega_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \omega^2 \end{pmatrix} \Omega_1\, R_{13}(\phi_m) \quad \text{and} \quad \phi_m = \frac{\pi m}{n} \;, \tag{47}$$

$$R_{12}(\theta) = \begin{pmatrix} \cos\theta & \sin\theta & 0 \\ -\sin\theta & \cos\theta & 0 \\ 0 & 0 & 1 \end{pmatrix} \tag{48}$$

and P_1 as defined in (43). The CP invariants are

$$\begin{aligned} J_{CP,3b1} &= -\frac{1}{6\sqrt{6}}\, \sin 3\,\phi_m \sin 3\,\phi_s \sin 2\theta \;\;, \\ I_{1,3b1} &= \frac{4}{9}\,(-1)^{k_2+1}\, \sin^2\phi_m \sin 3\,\phi_s \sin\theta \,\left(\cos 3\,\phi_s \sin\theta - \sqrt{2}\cos\phi_m \cos\theta\right) \;\;, \\ I_{2,3b1} &= \frac{4}{9}\,(-1)^{k_1+k_2+1}\, \sin^2\phi_m \sin 3\,\phi_s \cos\theta \left(\cos 3\,\phi_s \cos\theta + \sqrt{2}\cos\phi_m \sin\theta\right) \end{aligned} \tag{49}$$

and

$$I_{3,3b1} = \frac{1}{9}\,(-1)^{k_1+1}\, \cos\phi_m\, \sin 3\,\phi_s\, \left(4\cos\phi_m \cos 3\,\phi_s \cos 2\theta + \sqrt{2}\, \cos 2\,\phi_m\, \sin 2\theta\right) \;. \tag{50}$$

The expressions of the CP invariants in terms of ϕ_s and θ are considerably simplified in the case $m = n/2$ ($\phi_m = \pi/2$). And thus we can easily extract the Majorana and Dirac phases

$$\sin\alpha = (-1)^{k_2+1}\, \sin 6\,\phi_s \;\;,\;\; \sin\beta = (-1)^{k_1+k_2+1}\, \sin 6\,\phi_s \;\;,\;\; \sin\delta \approx \pm \sin 3\,\phi_s \;. \tag{51}$$

The last approximation holds, because $\theta \approx \pi/2$ (see [27] for details), and the plus (minus) sign refers to θ smaller (larger) than $\pi/2$. For $s = n/2$, $\sin\delta$ is actually independent of θ [27] and the approximation in (51) is exact. So, we find in this case $\sin\delta = \pm 1$, i.e. the Dirac is maximal, while the Majorana phases α and β are trivial for $s = n/2$.

For case 3 a) we consider instead of $U_{PMNS,3b1} = U_{R,3b1}$ in (46) the matrix without the permutation P_1

$$U_{PMNS,3a} = U_{R,3a} = \Omega_3\, R_{12}(\theta)\, K_\nu \;. \tag{52}$$

We find the same result for the Jarlskog invariant derived from $U_{PMNS,3b1}$ and $U_{PMNS,3a}$, while the CP invariants I_i are permuted among each other

$$\begin{aligned} & J_{CP,3a} = J_{CP,3b1} \ , \\ & I_{1,3a} = I_{3,3b1} \ , \ \ I_{2,3a} = -I_{1,3b1} \ \ \text{and} \ \ \ I_{3,3a} = -I_{2,3b1} \ . \end{aligned} \tag{53}$$

Obviously, the results for the mixing angles in case 3 a) are different from those obtained in case 3 b.1). Thus, the sets of parameters n, m, s and θ that lead to mixing angles in accordance with the experimental data are different in the two cases, see the extensive analysis in [27] for details. We remark that the formulae of mixing angles and CP invariants in case 3 b.1) and case 3 a) have certain symmetry properties. Applying the set of transformations

$$s \ \ \rightarrow \ \ n - s \ \ (\phi_s \ \rightarrow \ \pi - \phi_s) \ \ \text{and} \ \ \ \theta \ \ \rightarrow \ \ \pi - \theta \tag{54}$$

the mixing angles remain invariant, whereas the CP invariants change sign and thus also the sines of all three CP phases. For $s = n/2$, thus, sending θ into $\pi - \theta$ leaves the mixing angles untouched, while all CP phases change their sign. Using instead the transformations

$$m \ \ \rightarrow \ \ n - m \ \ (\phi_m \ \rightarrow \ \pi - \phi_m) \ \ \text{and} \ \ \ \theta \ \ \rightarrow \ \ \pi - \theta \tag{55}$$

the solar and reactor mixing angle as well as the CP invariants I_i remain unchanged, while $\sin^2\theta_{23}$ becomes $\cos^2\theta_{23}$ and J_{CP} changes sign. For more symmetry transformations of this type see table 6 in [27]. In the particular case $m = n/2$, we can conclude that the replacement of θ with $\pi - \theta$ does not affect the solar and reactor mixing angle as well as the CP invariants I_i, while the atmospheric mixing angle changes its octant and J_{CP} its sign. If we set $m = n/2$ and $s = n/2$ and apply the transformation $\theta \ \rightarrow \ \pi - \theta$, we see from (54) and (55) that the solar and reactor mixing angle are unchanged, the atmospheric mixing angle must be maximal, the CP invariants I_i must vanish and J_{CP} changes sign and is in general non-zero.

As in case 2), mixing matrices arising from permutations of the rows of the PMNS mixing matrices derived from $U_{R,3b1}$ and $U_{R,3a}$ do also admit good agreement with the experimental data on lepton mixing angles for certain choices of the parameters. The formulae for mixing angles and CP invariants can be obtained from those found for the PMNS mixing matrices with non-permuted rows by taking into account shifts in the parameters m and θ. Again, details can be found in [27].

3 Leptogenesis

We first collect several pieces of information regarding leptogenesis in general. We continue with the discussion of leptogenesis in our framework assuming a generic form of the PMNS mixing matrix. Subsequently, we turn to the detailed analysis of leptogenesis for the different scenarios of mixing, case 1) through case 3 b.1) and case 3 a). In doing so, we separate our discussion in an analytical and a numerical part. In order to not expand the latter too much we will concentrate on case 3 b.1) when studying the third class of mixing.

3.1 Preliminaries

As already mentioned, we focus on unflavored thermal leptogenesis as mechanism to generate correctly the measured value of the baryon asymmetry of the Universe $Y_B = (8.65 \pm 0.09) \times 10^{-11}$ [1]. Thus we assume RH neutrino masses to be larger than $10^{12}\,\text{GeV}$ [43]. Since the RH neutrino mass spectrum is not expected to be strongly hierarchical, we consider the contributions of all three RH neutrinos to the generation of the baryon asymmetry of the Universe

$$Y_B = \sum_{i=1}^{3} Y_{Bi} \ . \tag{56}$$

The quantities Y_{Bi} correspond to the part of the baryon asymmetry produced via the $i^{\rm th}$ RH neutrino. The latter can be expressed in the following way [44]

$$Y_{Bi} \approx 1.38 \times 10^{-3}\, \epsilon_i \sum_{j=1}^{3} \eta_{ij}\,. \tag{57}$$

The numerical value in (57) depends on the yield of RH neutrinos at high temperatures and on the sphaleron conversion factor, while ϵ_i is the CP asymmetry, generated in decays of the RH neutrino N_i, and η_{ij} is the efficiency factor which parametrizes the washout and decoherence effects of the lepton charge asymmetry generated in N_i decays due to lepton number violating interactions which involve the states N_j present in the thermal bath at temperatures $T \sim M_i$. The CP asymmetry ϵ_i, arising from the decay of the RH neutrino N_i with the mass M_i, is defined as

$$\epsilon_i = -\frac{\sum_\alpha [\Gamma(N_i \to H l_\alpha) - \Gamma(N_i \to H^\star \bar{l}_\alpha)]}{\sum_\alpha [\Gamma(N_i \to H l_\alpha) + \Gamma(N_i \to H^\star \bar{l}_\alpha)]} = -\frac{1}{8\pi} \sum_{j \neq i} \frac{{\rm Im}\left((\hat{Y}_D^\dagger \hat{Y}_D)^2_{ij}\right)}{(\hat{Y}_D^\dagger \hat{Y}_D)_{ii}}\, f(M_j/M_i)\,, \tag{58}$$

and the loop function $f(x)$ in the SM reads [45]

$$f(x) = x\left[\frac{1}{1-x^2} + 1 - (1+x^2)\ln\left(1+\frac{1}{x^2}\right)\right]\,. \tag{59}$$

Notice that the quantities ϵ_i are specified as the CP asymmetries for anti-leptons such that the sign of ϵ_i is the same as for Y_{Bi} in (57).

The matrix $\hat{Y}_D$ is given by the neutrino Yukawa coupling Y_D in the RH neutrino mass basis and thus reads with our conventions, see (5) and (11),

$$\hat{Y}_D = Y_D\, U_R\,. \tag{60}$$

The efficiency factors η_{ij} are expected to lie in the interval $10^{-3} \lesssim \eta_{ij} \lesssim 1$, see comments at the end of the next subsection and figure 2. Their computation requires in general the numerical solution of Boltzmann equations. However, in our scenario several instances allow for a simplified treatment. The three different RH neutrinos N_i couple at LO to orthogonal linear combinations of the lepton flavors, since we find that the Yukawa coupling $\hat{Y}_D$ is proportional to the unitary matrix U_R, if we take into account that Y_D is proportional to the identity matrix at LO, see (8). The RH neutrino masses are taken to be larger than $10^{12}\,{\rm GeV}$ so that lepton flavor dynamics is not resolved at the temperatures relevant for leptogenesis and the Boltzmann equations of the three orthogonal lepton charge asymmetries generated by the decays of each RH neutrino are almost independent [44]. In the RH neutrino mass basis the efficiency factors η_{ij} then reduce to

$$\eta_{ij} \approx \eta_{ii}\, \delta_{ij}\,. \tag{61}$$

Furthermore, we constrain RH neutrino masses to be smaller than $10^{14}\,{\rm GeV}$. Thus, possible washout effects due to scattering processes which violate lepton number by two units are out of equilibrium and the efficiency factors η_{ii} in (61) can be approximated well as [46]

$$\eta_{ii} = \left(\frac{3.3 \times 10^{-3}\,{\rm eV}}{\tilde{m}_i} + \left(\frac{\tilde{m}_i}{0.55 \times 10^{-3}\,{\rm eV}}\right)^{1.16}\right)^{-1} \tag{62}$$

with $\tilde{m}_i$ being defined as

$$\tilde{m}_i = v_{\rm EW}^2\, \frac{(\hat{Y}_D^\dagger \hat{Y}_D)_{ii}}{M_i}\,. \tag{63}$$

Using that the RH neutrino masses M_i are directly related to the light neutrino masses m_i at LO, see (16), and are, in particular, not degenerate, we can show that the relative mass splitting of any pair N_i and N_j $(i \neq j)$ satisfies the bound

$$\frac{|M_i - M_j|}{M_i} \;\approx\; \left|1 - \frac{m_i}{m_j}\right| \gg \frac{\left|(\hat{Y}_D^\dagger \hat{Y}_D)_{ij}\right|}{16\,\pi^2} \;\approx 3 \times 10^{-(6 \div 7)} \left(\frac{\kappa}{10^{-3}}\right)\,, \tag{64}$$

for y_0 given in (69) and typical values of the expansion parameter κ, see (79). Thus, the perturbative expansion in Y_D of the CP asymmetries in (58) is still valid, see, e.g., [47, 48]. Similarly, thanks to the relation in (16) we can express the argument of the loop function $f(x)$ in (58) in terms of m_i

$$f(M_j/M_i) \approx f(m_i/m_j) \; . \tag{65}$$

In addition, using that Y_D is proportional to the identity matrix at LO, see (8), and that (16) holds, we find

$$\tilde{m}_i \approx m_i \; . \tag{66}$$

Thus, also the efficiency factors η_{ii} are functions of the light neutrino masses

$$\eta_{ii} \approx \eta(m_i) \; . \tag{67}$$

We can distinguish two regimes: for $\tilde{m}_i \approx m_i \lesssim 1.1 \times 10^{-3}\,\text{eV}$ we are in the weak washout regime, while for larger values of m_i in the strong washout regime, see [49]. In the first case the neutrino Yukawa interactions are so small that the number density of RH neutrinos does not reach thermal equilibrium. Consequently, there is a partial cancellation between the anti-lepton asymmetry generated during the production of the RH neutrinos and the lepton asymmetry produced by their subsequent decays. In the second case the abundance of RH neutrinos matches the equilibrium density and a sufficiently large lepton asymmetry can be entirely realized from the out-of-equilibrium decays of RH neutrinos.

Given the strong correlation between light and RH neutrino masses, focussing on the range $10^{12}\,\text{GeV} \lesssim M_i \lesssim 10^{14}\,\text{GeV}$ also constrains the masses m_i. In particular, we use as range for the lightest neutrino mass m_0 in our numerical analysis

$$5 \times 10^{-4}\,\text{eV} \lesssim m_0 \lesssim 0.1\,\text{eV} \; . \tag{68}$$

This corresponds, if we fix the mass of the heaviest RH neutrino to $10^{14}\,\text{GeV}$, to the following interval of the Yukawa coupling y_0

$$0.04 \lesssim \; y_0 \; \lesssim 0.6 \; . \tag{69}$$

In this case automatically the masses of the two lighter RH neutrinos lie in the interval $10^{12}\,\text{GeV} \lesssim M_i \lesssim 10^{14}\,\text{GeV}$. The coupling y_0 is (slightly) smaller than an order one number.[6]

As has been noticed in [32], if RH neutrinos transform in an irreducible three-dimensional representation of a flavor group G_f, the CP asymmetries ϵ_i vanish in the limit of an exact residual symmetry in the neutrino sector, since the combination $\hat{Y}_D^\dagger \hat{Y}_D$ is always proportional to the identity matrix. This also happens in our scenario. For this reason, corrections δY_D to the Yukawa coupling Y_D of the neutrinos play a crucial role for having successful leptogenesis. We can expand the matrix $\hat{Y}_D^\dagger \hat{Y}_D$ which enters in the expression of the CP asymmetries (58) as

$$\hat{Y}_D^\dagger \hat{Y}_D \approx U_R^\dagger \left((Y_D^0)^\dagger Y_D^0 + (\delta Y_D)^\dagger Y_D^0 + (Y_D^0)^\dagger \delta Y_D \right) U_R \tag{70}$$

at lowest orders in the correction δY_D. The first term in this expansion is diagonal and therefore does not contribute to the numerator of ϵ_i. However, the other two terms lead to off-diagonal entries in the matrix combination $\hat{Y}_D^\dagger \hat{Y}_D$. This expansion in δY_D also corresponds to an expansion in the parameter κ, simply because any correction δU_R to the diagonalization matrix of the RH neutrino mass matrix can only be effective, if at the same time also the correction δY_D is present. Thus, we expect from (58) and (70)

$$\epsilon_i \propto \kappa^2 \; . \tag{71}$$

[6] Its suppressed value can be easily achieved by an additional Z_2 symmetry under which, for example, only RH neutrinos transform. Then the Yukawa coupling Y_D of the neutrinos, see (5), does not originate anymore from a (different) renormalizable operator(s), but requires an insertion of a field whose VEV spontaneously breaks the Z_2 symmetry. This idea has been applied in e.g. [34]. Note that in this case also the correction δY_D in (27) will not only be suppressed by κ, but also by the expansion parameter associated with the VEV of the Z_2 group breaking field.

This observation is in accordance with the results obtained in models with a flavor symmetry only [32–34].[7] Furthermore, using (58) and (70) we see that the leading term in the CP asymmetries is independent of the parameter y_0. Since this statement does not hold for higher order terms in κ, we study this issue in our numerical analysis (see details in subsection 3.3). We show that for y_0 in the range given in (69) and κ, $\tilde{\kappa}$, see definition in (74), chosen as in (79), the relative difference between the exact expression of the CP asymmetries defined in (58) and their LO expansion in κ is typically smaller than 0.1.

In order to maximize the predictability of the scenario and to reduce its model-dependence we would like to require that the CP asymmetries ϵ_i are independent of the phases contained in the correction term δY_D. Apart from the simple case of real corrections this also holds,[8] if

$$(\delta Y_D)^\dagger Y_D^0 \tag{72}$$

is a symmetric matrix, since in this case the two latter terms in (70) are complex conjugated to each other so that clearly the phases contained in δY_D are irrelevant at the lowest order. They appear, however, in the term neglected in (70). In our scenario Y_D^0 is proportional to the identity matrix, see (8), and thus δY_D must be a symmetric matrix itself in order to fulfil the constraint that the matrix in (72) is symmetric. For δY_D in (27) this is true and thus we immediately know that all our results for the CP asymmetries will be independent of the phases of the parameters $z_{1,2}$ at lowest order. In our numerical analysis we also study the effect of the latter phases, entering in the subdominant terms, on the results for the CP asymmetries. We compute the relative difference between the results using the expression of the CP asymmetries in (58), which include phases of the complex parameters $z_{1,2}$ in (27), and the corresponding LO expressions in κ and find it typically to be smaller than 0.1 in our numerical analysis, see discussion of case 1) in subsection 3.4.

Finally, we note that also in several models with the flavor symmetry A_4 only [32–34, 36] the LO results of the CP asymmetries ϵ_i turn out to be independent of the phases of the correction δY_D and to only depend on the phases appearing in the diagonalization matrix U_R that are identified with the Majorana phases at low energies. This result can be traced back to the fact that the matrix combination given in (72) is a symmetric matrix also in these models. If this condition is not fulfilled, the independence of the CP asymmetries of the phases of the correction δY_D can be insured, if δY_D is chosen to be real.

3.2 General results

Before discussing explicitly the results for the different cases, case 1) through case 3 b.1) and case 3 a), we would like to work out the form of the CP asymmetries obtained for $U_R = U_{PMNS}$. In this case mixing angles (and CP phases) are not assumed to be predicted by any flavor or CP symmetry, but only to have values compatible with the experimental data. Thus, we parametrize $U_R = U_{PMNS}$ using the convention given in (166) in appendix A.1. The form of the neutrino Yukawa coupling Y_D is taken to be the sum of the LO term Y_D^0 in (8) and the correction δY_D in (27).

As argued below (72), at the dominant order in κ only the real parts of the parameters $z_{1,2}$ (see (27)) enter the expressions of the CP asymmetries. Thus, we define

$$\mathrm{Re}(z_1) = z\,\cos\zeta \quad \text{and} \quad \mathrm{Re}(z_2) = z\,\sin\zeta\,. \tag{73}$$

The vanishing of one of the parameters z_1 and z_2 refers to particular choices of ζ, $\zeta = \pi/2, 3\pi/2$ and $\zeta = 0, \pi$, respectively. As explained in appendix D, in an explicit model such special values can be achieved, for example, with a particular alignment of the VEV of a flavor symmetry breaking field. Since κ is always accompanied by z, we furthermore define as (effective) expansion parameter

$$\tilde{\kappa} = \kappa\, z\,. \tag{74}$$

[7] Notice that the quadratic dependence on the expansion parameter originates from generating the lepton asymmetry via unflavored leptogenesis.

[8] The reality of δY_D could be ensured with the help of a CP symmetry. In our present scenario this is not necessary, but would in any case require that this CP symmetry is distinct from the one preserved in the neutrino sector, since otherwise the CP asymmetries ϵ_i would vanish.

The analytic expressions of the CP asymmetries at LO in $\tilde{\kappa}$ in terms of the mixing angles, CP phases and ζ are in general rather lengthy. However, they considerably simplify, if we choose $z_2 = 0$ or equivalently $\zeta = 0, \pi$,

$$\begin{aligned}
\epsilon_1 &\approx -\frac{3\,\tilde{\kappa}^2}{2\,\pi}\,\cos^2\theta_{12}\,\cos^2\theta_{13}\left[\cos^2\theta_{13}\,\sin^2\theta_{12}\,\sin\alpha\,f\left(\frac{m_1}{m_2}\right) + \sin^2\theta_{13}\,\sin\beta\,f\left(\frac{m_1}{m_3}\right)\right] \\
&= -\frac{3\,\tilde{\kappa}^2}{2\,\pi}\left[I_1\,f\left(\frac{m_1}{m_2}\right) + I_2\,f\left(\frac{m_1}{m_3}\right)\right] \qquad (75) \\
\epsilon_2 &\approx \frac{3\,\tilde{\kappa}^2}{2\,\pi}\,\sin^2\theta_{12}\,\cos^2\theta_{13}\left[\cos^2\theta_{12}\,\cos^2\theta_{13}\,\sin\alpha\,f\left(\frac{m_2}{m_1}\right) - \sin^2\theta_{13}\,\sin\left(\beta-\alpha\right)\,f\left(\frac{m_2}{m_3}\right)\right] \\
&= \frac{3\,\tilde{\kappa}^2}{2\,\pi}\left[I_1\,f\left(\frac{m_2}{m_1}\right) - I_3\,f\left(\frac{m_2}{m_3}\right)\right] \qquad (76) \\
\epsilon_3 &\approx \frac{3\,\tilde{\kappa}^2}{8\,\pi}\,\sin^2 2\theta_{13}\left[\cos^2\theta_{12}\,\sin\beta\,f\left(\frac{m_3}{m_1}\right) + \sin^2\theta_{12}\,\sin\left(\beta-\alpha\right)\,f\left(\frac{m_3}{m_2}\right)\right] \\
&= \frac{3\,\tilde{\kappa}^2}{2\,\pi}\left[I_2\,f\left(\frac{m_3}{m_1}\right) + I_3\,f\left(\frac{m_3}{m_2}\right)\right]. \qquad (77)
\end{aligned}$$

The first line of these equations tells us that the sign of all CP asymmetries depends on the sines of the Majorana phases α and β as well as a possible sign coming from the loop function $f(m_i/m_j)$, $i \neq j$. Especially, the lepton mixing angles θ_{13} and θ_{12} do not have any influence on the sign. As we see, the CP asymmetries ϵ_i do neither depend on the Dirac phase δ nor on the atmospheric mixing angle θ_{23}. In the second line of these equations we have used the CP invariants $I_{1,2,3}$ given in appendix A.1. This establishes the connection between low and high energy phases.

In order to understand the sign of the CP asymmetries and to estimate the size of the expansion parameter $\tilde{\kappa}$ necessary in order to achieve ϵ_i that permits a sufficiently large baryon asymmetry of the Universe, we first briefly analyze the behavior of the loop function. The information on the light neutrino mass spectrum, whether it follows NO or IO, affects the value of the loop function $f(x)$. In particular, it affects also the sign of $f(x)$ as shown in figure 1, which displays the behavior of the loop function $f(x)$ for the six different arguments m_i/m_j, $i \neq j$, $i, j = 1, 2, 3$, with respect to the lightest neutrino mass m_0. In the panels on the left we assume light neutrinos with NO, while in the right ones they are supposed to follow IO. The solar and atmospheric mass squared differences, $\Delta m^2_{\rm sol}$ and $\Delta m^2_{\rm atm}$, take values in their $3\,\sigma$ range given in [14] and reported in appendix A.2, see (173) and (176). These ranges determine the width of the blue and red bands in the six plots of figure 1. This is particular relevant for a neutrino mass spectrum with IO, since m_1 and m_2 are almost degenerate in this case. If we assume a hierarchical light neutrino mass spectrum, namely for $m_0 \lesssim 0.004$ eV,[9] $f(m_i/m_j)$ is always negative, but $f(m_1/m_2)$ for IO which is large and positive. For large m_0, $f(m_i/m_j) \approx -f(m_j/m_i)$, $i \neq j$, holds showing that half of the loop functions attain positive and half negative values. Notice also within the whole range of m_0 shown in figure 1 the sign of the loop function does not change in the case of $f(m_2/m_1)$ and $f(m_3/m_{1,2})$ for NO and in the case of $f(m_1/m_{2,3})$ and $f(m_2/m_{1,3})$ for IO (all of them lead to negative sign apart from $f(m_1/m_2)$ for IO). Relevant is also the size of $f(x)$. As is well-known for IO $f(m_1/m_2) \approx -f(m_2/m_1)$ always and very large, thus enhancing both asymmetries ϵ_1 and ϵ_2, for example see figure 3 in subsection 3.3, whereas $f(m_{1,2}/m_3)$ and $f(m_3/m_{1,2})$ are usually subdominant for all values of m_0 and for both mass orderings. Thus, if no special cancellations occur, we expect the contribution from ϵ_3 to the baryon asymmetry of the Universe always to be subdominant and thus unlikely to determine the sign of the latter.

We also estimate the size of the CP invariants using the $3\,\sigma$ intervals of the reactor and the solar

[9] This value is chosen, since $f(m_1/m_2)$ changes its sign there for NO.



Figure 1: Behavior of the loop function $f(x)$ shown in (59) for the six different arguments m_i/m_j, $i \neq j$, $i, j = 1, 2, 3$, with respect to the lightest neutrino mass m_0. Plots on the left (right) side correspond to a light neutrino mass spectrum with NO (IO). Note that m_0 is given by m_1 in the former and by m_3 in the latter case. The mass squared differences $\Delta m^2_{\rm sol}$ and $\Delta m^2_{\rm atm}$ are taken to be in their experimentally preferred $3\,\sigma$ range given in [14] and reported in (173) and (176) in appendix A.2, respectively. This variation explains the width of the red and blue curves.

mixing angles shown in appendix A.2

$$I_1 \approx (0.19 \div 0.22)\,\sin\alpha\,,$$
$$I_2 \approx (1.2 \div 1.8)\,\times 10^{-2}\,\sin\beta \quad \text{and} \quad I_3 \approx (4.9 \div 8.4)\,\times 10^{-3}\,\sin(\beta-\alpha)\,. \tag{78}$$

Thus, if the Majorana phase α is not small and the value of $f(m_1/m_2)$ and $f(m_2/m_1)$ is not close to zero for a particular choice of the lightest neutrino mass m_0 (only occurring for NO and for $m_0 \approx 0.004$ eV), we expect the contributions involving I_1 ($\sin\alpha$) to dominate over the others. As a consequence, the absolute values of the CP asymmetries ϵ_1 and ϵ_2 should be (much) larger than $|\epsilon_3|$. This suppression of ϵ_3 (in addition to the one coming from the loop function $f(x)$) can also be understood by noticing that ϵ_3 is proportional to $\sin^2 2\theta_{13}$, whereas $\epsilon_{1,2}$ are proportional to $\cos^2\theta_{13}$. With (75) the expected size of the CP asymmetry is found to be: for z being of order one and no particular enhancement or suppression of the loop function, i.e. $f(x)$ is also of order one, the expansion parameter has to fulfill

$$\kappa\,,\ \tilde{\kappa} \gtrsim 10^{-3} \tag{79}$$

in order to achieve $|\epsilon_{1,2}| \gtrsim 10^{-6}$ which is the typical size of the CP asymmetry necessary to ensure successful leptogenesis [49]. Our estimate in (79) for κ is naturally reproduced in typical non-SUSY as well as SUSY models, see our sketch of models in appendix D.



Figure 2: Behavior of the efficiency factor $\eta(m_i)$ defined in (62) for the three different neutrino masses m_i with respect to the lightest neutrino mass m_0. Plots on the left (right) side correspond to a light neutrino mass spectrum with NO (IO). Note that m_0 is given by m_1 in the former and by m_3 in the latter case. The mass squared differences $\Delta m^2_{\rm sol}$ and $\Delta m^2_{\rm atm}$ are taken to be in their experimentally preferred $3\,\sigma$ ranges given in [14] and reported in (173) and (176) in appendix A.2, respectively. This variation explains the width of the curves.

In the other limiting case, $z_1 = 0$ and thus $\zeta = \pi/2, 3\,\pi/2$, one can show that all three CP phases contained in the PMNS mixing matrix enter the expression of the CP asymmetries. In order to simplify the form of the latter, we thus consider cases in which two of the three CP phases are trivial. We focus here on the case in which the Majorana phase β and the Dirac phase δ are trivial, i.e.

$$\beta = k_\beta\,\pi \;\;,\;\; \delta = k_\delta\,\pi \quad \text{with} \quad k_{\beta,\delta} = 0, 1\;, \tag{80}$$

since this also happens in case 1), see CP invariants in (34). We find compact formulae for $\epsilon_{1,3}$, if the conditions in (80) are imposed,

$$\begin{aligned}
\epsilon_1 &\approx -\frac{\tilde{\kappa}^2}{2\,\pi}\,\sin\alpha\left[\frac{1}{2}\,\left(1+\sin^2\theta_{13}\right)\,\sin 2\theta_{12}\,\cos 2\theta_{23} + (-1)^{k_\delta}\,\sin\theta_{13}\,\cos 2\theta_{12}\,\sin 2\theta_{23}\right]^2 f\left(\frac{m_1}{m_2}\right)\;, \\
\epsilon_3 &\approx \frac{\tilde{\kappa}^2}{2\,\pi}\,(-1)^{k_\beta+1}\,\sin\alpha\,\cos^2\theta_{13}\left[\cos\theta_{12}\,\sin 2\theta_{23} + (-1)^{k_\delta}\,\sin\theta_{13}\,\sin\theta_{12}\,\cos 2\theta_{23}\right]^2 f\left(\frac{m_3}{m_2}\right)
\end{aligned} \tag{81}$$

while ϵ_2 can be expressed in terms of the other two CP asymmetries

$$\epsilon_2 = -\epsilon_1\,f\left(\frac{m_2}{m_1}\right)\left(f\left(\frac{m_1}{m_2}\right)\right)^{-1} - \epsilon_3\,f\left(\frac{m_2}{m_3}\right)\left(f\left(\frac{m_3}{m_2}\right)\right)^{-1}\;. \tag{82}$$

This occurs, because $\mathrm{Im}\left((\hat{Y}_D^\dagger\hat{Y}_D)^2_{ij}\right)$ vanishes for $ij = 13, 31$ (not only at LO in $\tilde{\kappa}$, but exactly) in this case. At this order in the expansion parameter $\tilde{\kappa}$ the CP asymmetries ϵ_1 and ϵ_3 depend on three different pieces: the sine of the non-trivial Majorana phase α, the loop function $f(x)$ and a combination of trigonometric functions of the mixing angles that forms a square. Thus, the sign of ϵ_1 and ϵ_3 depends on the sign of $\sin\alpha$ (and whether $\beta = 0$ or π) and the sign of the loop function. Like in the case above, we reach the conclusion that the sign of the CP asymmetries can be fixed with the knowledge of the Majorana phase(s). Results very similar to those in (81-82) are obtained, if the Majorana phase α is trivial instead of β. For completeness, we mention the formulae belonging to this choice of phases in appendix C. Moreover, we can analyze the case in which both Majorana phases are trivial, see also appendix C. All three CP asymmetries ϵ_i are still non-zero in this case and are all proportional to $\sin\delta$. However, the sign of the CP asymmetries ϵ_i does not only depend on the sign of $\sin\delta$ (and the loop function $f(x)$) in this case, but also, for example, on the octant of the atmospheric mixing angle $\cos 2\theta_{23} \lessgtr 0$, see (185).

Before closing this general part we would like to also comment in a quantitative way on the efficiency factors $\eta_{ii} \approx \eta(m_i)$, given in (62), that are necessary for the computation of the baryon

asymmetry of the Universe. We show the variation of the efficiency factor with respect to the light neutrino masses for NO (left panel) and IO (right panel) in figure 2. Like in the case of the loop function, the width of the curves is given by the 3σ intervals of the solar and atmospheric mass squared differences. As we can see, the efficiency factor can reach sizable values, up to 0.2, in the weak washout regime, i.e. for $\tilde{m}_i \approx m_i \lesssim 1.1 \times 10^{-3}$ eV. Conversely, in case of strong washout (for $m_i > 1.1 \times 10^{-3}$ eV), $\eta(m_i)$ is suppressed by one or more orders of magnitude. In particular, for $m_i \gtrsim \sqrt{\Delta m^2_{\rm sol}} \approx 0.009$ eV ($m_i \gtrsim \sqrt{|\Delta m^2_{\rm atm}|} \approx 0.05$ eV) we find $\eta(m_i) \lesssim 0.04$ ($\eta(m_i) \lesssim 0.001$). In the case of NO spectrum, the maximum efficiency is obtained in the production of Y_{B_1}, since $\eta(m_1)$ is the largest, unless the CP asymmetry ϵ_1 is suppressed. The strongest washout effects are expected for Y_{B_3}, which is controlled by the heaviest neutrino mass m_3. The opposite happens in the case of a spectrum with IO, where the lightest neutrino mass is m_3, while $m_2 \approx m_1 \gtrsim \sqrt{|\Delta m^2_{\rm atm}|}$. This does not necessarily mean that $\epsilon_{1,2}$ become much suppressed for IO, since the strong suppression of the baryon asymmetry Y_{B_i} due to washout effects can be easily compensated by the large enhancement of the loop functions in the expression of the CP asymmetries, as shown in figure 1. The latter argument has also to be taken into account in case of a QD neutrino mass spectrum.

3.3 Analytical discussion

After the general analysis we discuss the features of case 1) to case 3 b.1) and case 3 a) in an analytical way first. Then, we also show a numerical study for examples of each of the cases.

Leptogenesis in case 1)

We turn now to the predictions of the baryon asymmetry of the Universe for case 1). Accordingly, we express the CP asymmetries, as defined in (58), in terms of the relevant parameters which characterize the lepton mixing and the phases of the PMNS mixing matrix in this case, namely θ and the group theoretical parameters n and s (in the combination ϕ_s) –see (31-33). The expressions at LO in the expansion parameter $\tilde{\kappa}$, given in (74), are

$$\begin{aligned}
\epsilon_1 &\approx (-1)^{k_1} \frac{\tilde{\kappa}^2}{3\,\pi} \cos^2(\theta+\zeta)\; f\left(\frac{m_1}{m_2}\right) \sin 6\,\phi_s \\
&= -\frac{3\,\tilde{\kappa}^2}{2\,\pi} I_1(\theta \to \theta+\zeta)\; f\left(\frac{m_1}{m_2}\right) \qquad (83)\\
\epsilon_2 &\approx (-1)^{k_1+1} \frac{\tilde{\kappa}^2}{3\,\pi} \left(\cos^2(\theta+\zeta)\; f\left(\frac{m_2}{m_1}\right) + (-1)^{k_2} \sin^2(\theta+\zeta)\; f\left(\frac{m_2}{m_3}\right)\right) \sin 6\,\phi_s \\
&= \frac{3\,\tilde{\kappa}^2}{2\,\pi} \left[I_1(\theta \to \theta+\zeta)\; f\left(\frac{m_2}{m_1}\right) - I_3(\theta \to \theta+\zeta)\; f\left(\frac{m_2}{m_3}\right)\right], \qquad (84)\\
\epsilon_3 &\approx (-1)^{k_1+k_2} \frac{\tilde{\kappa}^2}{3\,\pi} \sin^2(\theta+\zeta)\; f\left(\frac{m_3}{m_2}\right) \sin 6\,\phi_s \\
&= \frac{3\,\tilde{\kappa}^2}{2\,\pi} I_3(\theta \to \theta+\zeta)\; f\left(\frac{m_3}{m_2}\right). \qquad (85)
\end{aligned}$$

Here $I_{1,3}(\theta \to \theta+\zeta)$ have to be read as

$$I_1(\theta \to \theta+\zeta) = \frac{2}{9}(-1)^{k_1+1} \cos^2(\theta+\zeta) \sin 6\,\phi_s \quad \text{and} \quad I_3(\theta \to \theta+\zeta) = \frac{2}{9}(-1)^{k_1+k_2} \sin^2(\theta+\zeta) \sin 6\,\phi_s \qquad (86)$$

meaning that the free parameter θ is shifted by ζ, which characterizes the correction δY_D, see (73). Thus, the CP asymmetries ϵ_i can be formally written in terms of the CP invariants I_1 and I_3 given in (34) (I_2 is zero anyway). The expressions in (86) coincide with the CP invariants $I_{1,3}$, see (34), of case 1) for $\zeta = 0,\ \pi$ and thus represent a special case of the general one found in (75-77). We can also compare these formulae with those obtained in (81-82) for the general case. In doing so, we only have to remember that $\sin 6\,\phi_s$ can be identified with $\sin\alpha$, see (35). In particular, we see that

case 1) offers an example in which ϵ_1 and ϵ_3 only receive one contribution and ϵ_2 can be written in terms of the former two ones, see (82). With this explicit case we can also confirm several of the observations made in subsection 3.1, e.g. no explicit dependence on the Yukawa coupling y_0, see (8), at LO, the CP asymmetries are proportional to $\tilde{\kappa}^2$ and the phases of the parameters $z_{1,2}$ only enter at higher orders in $\tilde{\kappa}$, i.e. these terms are suppressed by $(\tilde{\kappa}/y_0)^\sigma$ with $\sigma \geq 1$ with the respect to the LO terms.

The sign of the CP asymmetries is only determined by ϕ_s and the one of the loop function $f(x)$ for a given value of m_0 and, in particular, is independent of the values of the parameters θ and ζ. The identification of $\sin 6\,\phi_s$ with $\sin\alpha$ also shows that in the case of no low energy CP violation, $s = 0$ or $s = n/2$, also no CP violation occurs at high energies. We note that replacing s by $n - s$ reverses the sign of all three CP asymmetries ϵ_i

$$\epsilon_i(n-s) = -\epsilon_i(s)\;. \tag{87}$$

This equality holds exactly. Furthermore, we note that at LO in $\tilde{\kappa}$ the CP asymmetries are periodic functions in ζ with the period π

$$\epsilon_i(\zeta) \;=\; \epsilon_i(\zeta \;+\; k\,\pi)\,, \quad \text{for } k = 0, 1, 2, \ldots \tag{88}$$

As can be checked by explicit computation, this does not hold, if we include the higher order terms in $\tilde{\kappa}$ in (83-85). Similarly, at LO in $\tilde{\kappa}$ the CP asymmetries ϵ_i remain invariant, if we replace θ by $\pi - \theta$ and ζ by $k\,\pi - \zeta$ where k is an integer chosen in such a way that $k\,\pi - \zeta$ lies in the fundamental interval $[0, 2\,\pi)$. Thus, we expect to obtain the very similar results (for a different value of the parameter ζ however) for θ in the two different admitted intervals, $0.168 \lesssim \theta \lesssim 0.195$ and $2.95 \lesssim \theta \lesssim 2.97$, see also table 1.

Leptogenesis in case 2)

In this case the parameters which determine lepton mixing angles and CP phases are ϕ_u and ϕ_v that characterize the chosen CP transformation X (remember u and v are related to s and t, see (38)) and the free parameter θ. Computing the CP asymmetries at the lowest order in $\tilde{\kappa}$ we obtain

$$\begin{aligned}
\epsilon_1 \;&\approx\; \frac{\tilde{\kappa}^2}{6\,\pi}\left[(-1)^{k_1} f\left(\frac{m_1}{m_2}\right)\Big(\left[\cos\left(\phi_u + 2\zeta\right) + \cos 2\theta\right]\sin\phi_v - \sin\left(\phi_u + 2\zeta\right)\sin 2\theta\cos\phi_v\Big)\right. \\
&\quad \left. +(-1)^{k_2+1} f\left(\frac{m_1}{m_3}\right)\sin 2\left(\phi_u - \zeta\right)\sin 2\theta\right] \\
&=\; -\frac{3\,\tilde{\kappa}^2}{2\,\pi}\left[I_1\left(\phi_u \to \phi_u + 2\,\zeta\right)\, f\left(\frac{m_1}{m_2}\right) + I_2\left(\phi_u \to \phi_u - \zeta\right)\, f\left(\frac{m_1}{m_3}\right)\right]\,,
\end{aligned} \tag{89}$$

$$\begin{aligned}
\epsilon_2 \;&\approx\; -\frac{\tilde{\kappa}^2}{6\,\pi}\left[(-1)^{k_1} f\left(\frac{m_2}{m_1}\right)\Big(\left[\cos\left(\phi_u + 2\zeta\right) + \cos 2\theta\right]\sin\phi_v - \sin\left(\phi_u + 2\zeta\right)\sin 2\theta\cos\phi_v\Big)\right. \\
&\quad \left. +(-1)^{k_1+k_2} f\left(\frac{m_2}{m_3}\right)\Big(\left[\cos\left(\phi_u + 2\zeta\right) - \cos 2\theta\right]\sin\phi_v + \sin\left(\phi_u + 2\zeta\right)\sin 2\theta\cos\phi_v\Big)\right] \\
&=\; \frac{3\,\tilde{\kappa}^2}{2\,\pi}\left[I_1\left(\phi_u \to \phi_u + 2\,\zeta\right)\, f\left(\frac{m_2}{m_1}\right) - I_3\left(\phi_u \to \phi_u + 2\,\zeta\right)\, f\left(\frac{m_2}{m_3}\right)\right]\,,
\end{aligned} \tag{90}$$

$$\begin{aligned}
\epsilon_3 \;&\approx\; \frac{\tilde{\kappa}^2}{6\,\pi}\left[(-1)^{k_2} f\left(\frac{m_3}{m_1}\right)\sin 2\left(\phi_u - \zeta\right)\sin 2\theta\right. \\
&\quad \left. +(-1)^{k_1+k_2} f\left(\frac{m_3}{m_2}\right)\Big(\left[\cos\left(\phi_u + 2\zeta\right) - \cos 2\theta\right]\sin\phi_v + \sin\left(\phi_u + 2\zeta\right)\sin 2\theta\cos\phi_v\Big)\right] \\
&=\; \frac{3\,\tilde{\kappa}^2}{2\,\pi}\left[I_2\left(\phi_u \to \phi_u - \zeta\right)\, f\left(\frac{m_3}{m_1}\right) + I_3\left(\phi_u \to \phi_u + 2\,\zeta\right)\, f\left(\frac{m_3}{m_2}\right)\right]\,.
\end{aligned} \tag{91}$$

Similar to case 1), we can express the LO results for the CP asymmetries in terms of the quantities I_i, see (39-40), if we shift the group theoretical parameter ϕ_u by appropriate multiples of ζ. Again, only if the latter is taken to be 0 or π, we re-cover expressions that depend on the CP invariants I_i, as defined in (39-40), and thus on the sines of the Majorana phases α and β, while the Dirac phase δ does not appear. In this special case we can match to the formulae found in (75-77) where the PMNS mixing matrix is of a general form. We also find in this case that ϵ_i fulfill at LO the equality in (88), i.e. the expressions are invariant under the shift of ζ in $\zeta + k\,\pi$ where k is an integer. In contrast to case 1), it is not straightforward to determine the sign of ϵ_i just by looking at (89-91). However, given that θ is close to 0 or π (or in the case of a matrix of the form as in (42) $\theta \approx \pi/2$), the terms proportional to $\sin 2\theta$ are likely to be suppressed compared to those proportional to $\cos(\phi_u + 2\zeta) \pm \cos 2\theta \approx \cos(\phi_u + 2\zeta) \pm 1$, unless $\sin\phi_v$ is small. An example for the latter case is $v = 0$ (if this choice of v is admitted) and we see that

$$\epsilon_i \propto (c_1 \sin(\phi_u + 2\zeta) + c_2 \sin 2(\phi_u - \zeta)) \sin 2\theta \tag{92}$$

with $c_{1,2}$ being expressions of the loop function $f(m_i/m_j)$ and $k_{1,2}$. This shows that the terms in ϵ_i are proportional in this case to $\sin(\phi_u + 2\zeta)$ or $\sin 2(\phi_u - \zeta)$ and thus for fixed u (ϕ_u) crucially depend on the choice of ζ; for example, changing the latter into $\zeta \pm \pi/2$ changes the overall sign of the CP asymmetries. This eventually explains why in our numerical analysis for $v = 0$ positive and negative values of the baryon asymmetry of the Universe Y_B are equally likely, see light-blue areas in figure 6. As a consequence, in this case a prediction of the sign of Y_B, depending on the low energy CP phases, is not possible. The expression in (92) also shows that the choice of the parameter θ becomes relevant, in particular whether $\theta \lesssim \pi/2$ or $\theta \gtrsim \pi/2$, and thus in this case changing the interval of θ will inevitably change the sign of the CP asymmetries. In general, however, if θ is replaced by $\pi - \theta$, our predictions of the CP asymmetries will only slightly differ, since the dominating terms in ϵ_i are proportional to $\sin\alpha$, see (89-91), and only $\sin\delta$ and $\sin\beta$ change sign.

We can also relate the expressions found in case 2) to those in case 1). As has been shown in [27], we can set $\theta = 0$ and identify the parameters of case 2) ϕ_u and v with $2\,\theta_1$ and $6\,s_1$ of case 1), respectively, assuming that the group index n is the same in both cases. Then, ϵ_i of case 2) coincide with those of case 1), in particular I_2 (for any argument) vanishes. Another way to relate case 2) and case 1) is to set $u = 0$ ($\phi_u = 0$) and to relate v with $6\,s$. The admitted intervals of θ of case 2) coincide with those of case 1), see table 1. Then, the results for the solar and reactor mixing angles are the same in both cases as well as the Majorana phase α, while the other phase β is trivial.[10] Consequently, for $\zeta = 0, \pi$ the CP asymmetries ϵ_i coincide in case 1) and case 2) for this choice of parameters. However, if $\zeta \neq 0, \pi$, we expect different results for both cases, since, in particular, the CP asymmetries in case 2) include additional terms proportional to $I_2\,(\phi_u \to \phi_u - \zeta)$.

We can also consider leptogenesis in the case in which the PMNS mixing matrix/the matrix U_R is given by one of the two matrices in (42). As explained in subsection 2.2, the formulae for mixing angles and CP phases can be derived from those given for case 2), i.e. for U_{PMNS}/U_R being as in (36) using the transformations displayed in (44) and (45), respectively. In the same vein, we can obtain the formulae for the CP asymmetries ϵ_i in the shifted cases. This is, in particular, relevant for our numerical discussion, since we consider indeed the case in which the permutation matrix P_1 is applied from the left to the matrix in (36). In the case we assume that the additional permutation originates from the Majorana mass matrix of the RH neutrinos and for this reason we have included it in the matrix U_R. However, in principle we can also consider a different situation in which U_R is still given by (36), whereas the PMNS mixing matrix is given by one of the matrices in (42). In such a situation the permutation arises from the charged lepton sector, since the charged lepton masses are non-canonically ordered. One may wonder whether this difference can lead to new results. Indeed, this is not the case and it is no restriction of our discussion to focus only on the scenario with the permutation being included in the RH neutrino sector. The other case can be simply obtained from the latter one by a permutation P of the (diagonal) elements of the correction

[10] The only differences lie in the values of the atmospheric mixing angle and the Dirac phase, since case 2) predicts both of them to be maximal, if u equals zero.

δY_D, i.e.

$$(\delta Y_D)_{P \text{ from } m_l} = P\,(\delta Y_D)_{P \text{ in } U_R}\,P^T\,. \tag{93}$$

Thus, we only need to re-define the parameters z_1 and z_2 appropriately. Clearly, this affects (mainly) the explicit value of the parameter ζ corresponding to a certain choice of the ratio of the real parts of z_1 and z_2, but not the general conclusion on the size and the sign of the baryon asymmetry of the Universe, obtained in the different cases.

Leptogenesis in case 3)

We finally consider the predictions for the baryon asymmetry in the last class of mixing patterns, which are classified as case 3 b.1) and case 3 a). As we show, the CP asymmetries predicted in the two cases are closely related.

The analytic approximations of the CP asymmetries in case 3 b.1) are expressed at LO in $\tilde{\kappa}$ as a function of the group theoretical quantities ϕ_m, ϕ_s and the free parameter θ, that is

$$\begin{aligned}
\epsilon_1 &\approx \frac{\tilde{\kappa}^2}{12\,\pi}\left[(-1)^{k_2} f\left(\frac{m_1}{m_2}\right)\left(4\sin^2(\phi_m+\zeta)\sin^2\theta\sin 6\,\phi_s+\sqrt{2}\left[\cos 3\,\phi_m-\cos\left(\phi_m-2\zeta\right)\right]\sin 3\,\phi_s\sin 2\theta\right)\right. \\
&\left.+(-1)^{k_1+k_2} f\left(\frac{m_1}{m_3}\right)\left(4\sin^2(\phi_m+\zeta)\cos^2\theta\sin 6\,\phi_s-\sqrt{2}\left[\cos 3\,\phi_m-\cos\left(\phi_m-2\zeta\right)\right]\sin 3\,\phi_s\sin 2\theta\right)\right] \\
&= -\frac{3\,\tilde{\kappa}^2}{2\,\pi}\left[\left(I_1\left(\phi_m\to\phi_m+\zeta\right)+(-1)^{k_2+1}R_-(\zeta)\right) f\left(\frac{m_1}{m_2}\right)+\left(I_2\left(\phi_m\to\phi_m+\zeta\right)+(-1)^{k_1+k_2}R_-(\zeta)\right) f\left(\frac{m_1}{m_3}\right)\right]
\end{aligned} \tag{94}$$

$$\begin{aligned}
\epsilon_2 &\approx -\frac{\tilde{\kappa}^2}{12\,\pi}\left[(-1)^{k_2} f\left(\frac{m_2}{m_1}\right)\left(4\sin^2(\phi_m+\zeta)\sin^2\theta\sin 6\,\phi_s+\sqrt{2}\left[\cos 3\,\phi_m-\cos\left(\phi_m-2\zeta\right)\right]\sin 3\,\phi_s\sin 2\theta\right)\right. \\
&\left.+(-1)^{k_1+1} f\left(\frac{m_2}{m_3}\right)\left(4\cos^2(\phi_m+\zeta)\cos 2\theta\sin 6\,\phi_s+\sqrt{2}\left[\cos 3\,\phi_m+\cos\left(\phi_m-2\zeta\right)\right]\sin 3\,\phi_s\sin 2\theta\right)\right] \\
&= \frac{3\,\tilde{\kappa}^2}{2\,\pi}\left[\left(I_1\left(\phi_m\to\phi_m+\zeta\right)+(-1)^{k_2+1}R_-(\zeta)\right) f\left(\frac{m_2}{m_1}\right)-\left(I_3\left(\phi_m\to\phi_m+\zeta\right)+(-1)^{k_1}R_+(\zeta)\right) f\left(\frac{m_2}{m_3}\right)\right]
\end{aligned} \tag{95}$$

$$\begin{aligned}
\epsilon_3 &\approx -\frac{\tilde{\kappa}^2}{12\,\pi}\left[(-1)^{k_1+k_2} f\left(\frac{m_3}{m_1}\right)\left(4\sin^2(\phi_m+\zeta)\cos^2\theta\sin 6\,\phi_s-\sqrt{2}\left[\cos 3\,\phi_m-\cos\left(\phi_m-2\zeta\right)\right]\sin 3\,\phi_s\sin 2\theta\right)\right. \\
&\left.+(-1)^{k_1} f\left(\frac{m_3}{m_2}\right)\left(4\cos^2(\phi_m+\zeta)\cos 2\theta\sin 6\,\phi_s+\sqrt{2}\left[\cos 3\,\phi_m+\cos\left(\phi_m-2\zeta\right)\right]\sin 3\,\phi_s\sin 2\theta\right)\right] \\
&= \frac{3\,\tilde{\kappa}^2}{2\,\pi}\left[\left(I_2\left(\phi_m\to\phi_m+\zeta\right)+(-1)^{k_1+k_2}R_-(\zeta)\right) f\left(\frac{m_3}{m_1}\right)+\left(I_3\left(\phi_m\to\phi_m+\zeta\right)+(-1)^{k_1}R_+(\zeta)\right) f\left(\frac{m_3}{m_2}\right)\right]
\end{aligned} \tag{96}$$

Like in the other cases, we have arranged the LO analytic approximations of ϵ_i in terms of the expressions of the CP invariants I_1, I_2 and I_3 for case 3 b.1), defined in (49-50). This requires that the group theoretical parameter ϕ_m is shifted by multiples of the parameter ζ as well as that we have to include an additional piece, parametrized as follows

$$R_\pm(\zeta) = \frac{\sqrt{2}}{9}\sin\frac{3\,\zeta}{2}\left[\sin\left(\phi_m-\frac{\zeta}{2}\right)\pm\sin 3\left(\phi_m+\frac{\zeta}{2}\right)\right]\sin 3\,\phi_s\sin 2\theta\,. \tag{97}$$

Like in the cases above we confirm all statements made in the general part, regarding the expansion in $\tilde{\kappa}$, the appearance of the Yukawa coupling y_0 and the dependence on the imaginary parts of the parameters $z_{1,2}$ which is only subdominant. We can furthermore check that the LO expressions of the CP asymmetries (94-96) are periodical in ζ with periodicity π. As in the other cases, this is only correct at LO and not at higher orders in $\tilde{\kappa}$. As in the general case, see (75-77), we find

that for $\zeta = 0, \pi$ the CP asymmetries can be written in terms of the CP invariants I_i in (49-50). This also shows that there is no explicit dependence of the Dirac phase in this case. As shown in [27], in case 3 b.1) the replacement of θ with $\pi - \theta$ does not give rise to a symmetry of the formulae in general, only if we replace at the same time m with $n - m$ or s with $n - s$, see table 6 in [27]. Thus, we only discuss the case a possible second interval for θ, if $m = n/2$, which is fulfilled in our numerical example, see subsection 3.4. In this case we see that the sines of the Majorana phases remain invariant, whereas the sine of the Dirac phase changes sign. Consequently, the CP asymmetries ϵ_i are similar to those obtained for θ in the first interval, unless $\sin\alpha$ is small and a prediction of the sign of ϵ_i becomes anyway impossible, see subsection 3.4. In the general case, we can make the following observation, deduced from the transformations in table 6 in [27]

$$\epsilon_i\,(n-s, \pi-\theta) = -\epsilon_i\,(s,\theta)\ . \tag{98}$$

This equality holds for all three asymmetries, at all orders in $\tilde{\kappa}$ and for all choices of the parameters $z_{1,2}$. We use this fact in our numerical analysis where we only display results for choices $s \leq n/2$.

The expressions in (94-96) considerably simplify for $m = n/2$ and $s = n/2$ (assuming that n is even)

$$\epsilon_1 \approx \frac{\tilde{\kappa}^2}{6\sqrt{2}\,\pi}\,(-1)^{k_2}\left[f\left(\frac{m_1}{m_2}\right) + (-1)^{k_1+1} f\left(\frac{m_1}{m_3}\right)\right]\sin 2\zeta\,\sin 2\theta\,, \tag{99}$$

$$\epsilon_2 \approx -\frac{\tilde{\kappa}^2}{6\sqrt{2}\,\pi}\,(-1)^{k_2}\left[f\left(\frac{m_2}{m_1}\right) + (-1)^{k_1+k_2} f\left(\frac{m_2}{m_3}\right)\right]\sin 2\zeta\,\sin 2\theta\,, \tag{100}$$

$$\epsilon_3 \approx \frac{\tilde{\kappa}^2}{6\sqrt{2}\,\pi}\,(-1)^{k_1+k_2}\left[f\left(\frac{m_3}{m_1}\right) + (-1)^{k_2} f\left(\frac{m_3}{m_2}\right)\right]\sin 2\zeta\,\sin 2\theta\,. \tag{101}$$

In this case it is clear that we cannot predict the sign of the CP asymmetries ϵ_i, since it crucially depends on the choice of the free parameter ζ. It also crucially depends on whether θ is smaller or larger than $\pi/2$. Thus, changing the admitted interval of θ, now two exist in general, leads to a change in the sign of ϵ_i. This observation is intimately related to the fact that both Majorana phases are trivial in this case, while the Dirac phase is maximal and its sign depends on θ, see (51). This is very similar to what we have observed in case 2), see discussion around (92). For $\zeta = 0, \pi$ the CP asymmetries are exactly zero at all orders in $\tilde{\kappa}$, while they take maximal (positive and negative) values for $\zeta = \pi/4$ or $3\pi/4$. For the special choices $\zeta = \pi/2, 3\pi/2$ the CP asymmetries at LO vanish as well. This observation is consistent with the fact that the formulae derived in the case in which the mixing matrix U_R is parametrized as general matrix and $\zeta = \pi/2, 3\pi/2$ lead also to zero CP asymmetries for maximal Dirac phase and maximal atmospheric mixing, see (183-185) in appendix C.

The expressions of ϵ_i corresponding to case 3 a) can be easily obtained from (94-96), using the relations among the CP invariants in the two cases, reported in (53), and the replacements

$$I_{1,3b1}\,(\phi_m \to \phi_m + \zeta) + (-1)^{k_2+1} R_-(\zeta) \to I_{1,3a}\,(\phi_m \to \phi_m + \zeta) + (-1)^{k_1} R_+(\zeta)\,, \tag{102}$$

$$I_{2,3b1}\,(\phi_m \to \phi_m + \zeta) + (-1)^{k_1+k_2} R_-(\zeta) \to I_{2,3a}\,(\phi_m \to \phi_m + \zeta) + (-1)^{k_2} R_-(\zeta)\,, \tag{103}$$

$$I_{3,3b1}\,(\phi_m \to \phi_m + \zeta) + (-1)^{k_1} R_+(\zeta) \to I_{3,3a}\,(\phi_m \to \phi_m + \zeta) + (-1)^{k_1+k_2+1} R_-(\zeta)\,. \tag{104}$$

If we consider instead U_{PMNS}/U_R with rows permuted, the given formulae for the CP asymmetries ϵ_i in (94-96) and (102-104) can still be applied, as long as we use the same replacements of the parameters m and θ that need to be taken into account when computing the mixing angles and CP phases, see end of subsection 2.2.

3.4 Numerical discussion

We first summarize our specific choices of several of the involved parameters that we use throughout the numerical discussion of the different cases and then we show results for examples of case 1),

case 2) as well as case 3 b.1). Eventually, we also comment on the results expected in case 3 a). We note that we always use the exact expressions for the CP asymmetries ϵ_i in our numerical analysis, i.e. those that are not expanded in the parameter $\tilde{\kappa}$ and are computed using (58) and the corresponding form of the mixing matrix $U_{R,i}$, see (31, 36, 42, 46) (and also implicitly (52)).

Preliminaries

The mixing matrix U_R contains in all cases the two parameters k_1 and k_2 in the matrix K_ν, see (13). In the following, we set, if not stated otherwise,

$$k_1 = 0 \quad \text{and} \quad k_2 = 0 \,. \tag{105}$$

Furthermore, we fix the heaviest RH neutrino mass to

$$M_{1(3)} = 10^{14}\,\text{GeV}\,, \quad \text{for NO (IO)}\,. \tag{106}$$

Using (16) we can then express y_0 as

$$y_0 \approx 1.82\,\sqrt{\frac{m_0}{1\text{ eV}}}\,. \tag{107}$$

The lightest neutrino mass m_0 is chosen in the interval in (68) and y_0, consequently, falls into the range in (69). The other neutrino masses (light and heavy ones) are then determined by the two experimentally measured mass squared differences, see (19-21). In particular, the masses of all three RH neutrinos are between $10^{12}\,\text{GeV}$ and $10^{14}\,\text{GeV}$ in this setting.

The expansion parameter $\tilde{\kappa}$ is taken as constant and fixed to

$$\tilde{\kappa} = 4 \times 10^{-3}\,. \tag{108}$$

This value is appropriate in order to achieve sizable CP asymmetries, since it satisfies the bound in (79). The parameter ζ that defines the relative size among the real parts of the two parameters z_1 and z_2, see (73), is either fixed to specific values

$$\zeta = 0,\,\pi \quad \text{equivalent to} \quad \text{Re}(z_2) = 0 \quad \text{or} \quad \zeta = \frac{\pi}{2},\,\frac{3\,\pi}{2} \quad \text{equivalent to} \quad \text{Re}(z_1) = 0 \tag{109}$$

or is taken to lie in the intervals $\mathcal{I}_i$ which is equivalent to constraining the real parts of $z_{1,2}$ to assume values in $[-2,-1/2]$ and/or $[1/2,2]$

$$\begin{array}{ll} \mathcal{I}_1 = [0.24, 1.33] & \text{equivalent to} \quad \text{Re}(z_1) > 0 \;\text{ and }\; \text{Re}(z_2) > 0\,, \\ \mathcal{I}_2 = [1.82, 2.90] & \text{equivalent to} \quad \text{Re}(z_1) < 0 \;\text{ and }\; \text{Re}(z_2) > 0\,, \\ \mathcal{I}_3 = [3.39, 4.47] & \text{equivalent to} \quad \text{Re}(z_1) < 0 \;\text{ and }\; \text{Re}(z_2) < 0\,, \\ \mathcal{I}_4 = [4.96, 6.04] & \text{equivalent to} \quad \text{Re}(z_1) > 0 \;\text{ and }\; \text{Re}(z_2) < 0\,. \end{array} \tag{110}$$

In our numerical discussion these different choices of the parameter ζ are indicated in different colors in the figures

$$\zeta = 0,\,\pi \;\text{ in red }\;,\;\; \zeta = \frac{\pi}{2},\,\frac{3\,\pi}{2} \;\text{ in green and }\; \zeta \in \mathcal{I}_i \;\text{ in the light-blue areas.} \tag{111}$$

For the numerical example in case 1) we not only show figures of the baryon asymmetry Y_B of the Universe, but also for the CP asymmetries ϵ_i. Moreover, we discuss in this example results for both the two possible light neutrino mass spectra, with NO and with IO. We also show in case 1) the effect of choosing the two parameters k_1 and k_2 differently from our standard choice in (105). Discussing subsequently an example for case 2) and one for case 3 b.1) we only focus a NO light neutrino mass spectrum as well as on the choice $k_1 = k_2 = 0$, since from the discussion in case 1) the relations to and changes coming from the other possible choices become apparent.

n	4			
θ	$0.168 \div 0.195$			
$\sin^2\theta_{12}$	$0.340 \div 0.342$			
$\sin^2\theta_{13}$	$0.0188 \div 0.0251$			
$\sin^2\theta_{23}$	$0.389 \div 0.403$			
$\sin\beta$	0			
$\sin\delta$	0			
s	0	1	2	3
$\sin\alpha$	0	1	0	-1

Table 1: **Case 1)**. Results for lepton mixing angles and sines of the CP phases α, β and δ. The former are within the experimentally preferred $3\,\sigma$ ranges. The choice $n = 4$ is the minimal value of n that can give rise to a non-trivial Majorana phase α, if we set $s = 1$ or $s = 3$. The sign of $\sin\alpha$ depends on k_1 and corresponds in this table to the choice $k_1 = 0$ (we also take $k_2 = 0$), compare to (35). An atmospheric mixing angle in the second octant, $0.597 \lesssim \sin^2\theta_{23} \lesssim 0.612$, is obtained for $2.95 \lesssim \theta \lesssim 2.97$ or if the second and third rows of the PMNS mixing matrix (and thus U_R) are exchanged (see [27] for details). Note that the results for the other two mixing angles and the CP phases are not altered.

Leptogenesis in case 1)

For all (even) n and all CP transformations with $s = 0, \ldots, n-1$, the values of the free parameter θ, which allows to reproduce the lepton mixing angles in accordance with the experimental data at the $3\,\sigma$ level or better, lie in the two intervals $0.168 \lesssim \theta \lesssim 0.195$ and $2.95 \lesssim \theta \lesssim 2.97$ [27]. These two choices are distinguished by the resulting value of the atmospheric mixing angle which lies in the first and second octant, respectively. In the following numerical analysis we focus on values of θ in the first interval. Using that at LO in $\tilde{\kappa}$ the expressions of the CP asymmetries remain invariant, if we exchange θ with $\pi - \theta$ and ζ with $k\,\pi - \zeta$ (k are suitable integers for keeping ζ in its fundamental interval between 0 and $2\,\pi$), we know that the results for θ in the second interval will be very similar to those for θ in the first one, only obtained for a different value of the free parameter ζ and up to corrections that are of order $\tilde{\kappa}^3$ at most. The minimal choice of n for which a non-trivial Majorana phase α can arise is $n = 4$, if s is taken to be odd. Then, α is maximal and $\sin\alpha$ is positive (negative) for $s = 1$ ($s = 3$). If s is even instead, we find no CP violation at all and consequently also vanishing CP asymmetries ϵ_i. Thus, this case is not considered any further. This information together with the results for the other mixing parameters is gathered in table 1.

For $n = 4$ and $s = 1$ (meaning $\alpha = \pi/2$) we show the predictions of the CP asymmetries ϵ_i in figure 3 as a function of the lightest neutrino mass m_0, in the case of a light neutrino mass spectrum with NO (left panel) and IO (right panel). We vary the parameter θ in the (first) interval $0.168 \lesssim \theta \lesssim 0.195$ and the solar and atmospheric mass squared differences within their 3σ ranges, determined by the global fit analysis [14], see appendix A.2. First of all, we notice that the CP asymmetry ϵ_1, generated in N_1 decays, is suppressed for $\zeta \approx \pi/2$ and $3\pi/2$, see green area in the upper panels of figure 3, since ϵ_1 is proportional to $\cos^2(\theta+\zeta)$ at LO in $\tilde{\kappa}$, see (83), and θ is small (or close to π). It does not exactly vanish at LO, since, indeed, an exact cancellation can only occur, if $\theta+\zeta$ equals $\pi/2$ or $3\pi/2$, which explains the light-blue region giving rise to values of ϵ_1 even smaller than those shown in the green area. Conversely, the absolute value of ϵ_1 is maximized for $\zeta \approx 0$ and π, as can be read off from the red area in the upper panels of figure 3. Similarly, ϵ_3 is suppressed for $\zeta \approx 0$ and π, compare red area in the lower panels of figure 3, because it is proportional to $\sin^2(\theta+\zeta)$ at LO in $\tilde{\kappa}$, see (85), while this proportionality leads to an enhancement of the CP asymmetry ϵ_3 for $\zeta \approx \pi/2$ and $3\pi/2$. Since θ is small (close to π), but not zero (exactly π), $\epsilon_{1(3)}$ do not vanish, if ζ is chosen as $\pi/2$, $3\pi/2$ (0, π), as can be seen best in the left upper plot of figure 3. The behavior of ϵ_2 is more complex, if light neutrinos follow NO, since it has two different contributions that can be of



Figure 3: **Case 1).** CP asymmetries for $n = 4$, $s = 1$, $k_1 = k_2 = 0$ and $\tilde{\kappa} = 0.004$ as function of the lightest neutrino mass m_0 for NO (left panel) and IO (right panel). The Majorana phase in this case is $\alpha = \pi/2$. The parameter θ is taken in the interval $0.168 \lesssim \theta \lesssim 0.195$. The neutrino mass squared differences are varied in their $3\,\sigma$ ranges [14]. Red and green areas correspond to the special choices $\zeta = 0,\ \pi$ and $\zeta = \pi/2,\ 3\pi/2$, respectively. The light-blue area is obtained for ζ lying in one of the four intervals $\mathcal{I}_i$, reported in (110). Note the different scale of the vertical axis chosen in the plots for ϵ_1 and ϵ_2 and IO, showing the enhancement of the CP asymmetries that arises from the (near) degeneracy of m_1 and m_2, if neutrinos follow IO (compare also the behavior of the loop function $f(x)$ in figure 1).

similar size, see (84). If m_0 is small, i.e. $m_0 \lesssim 0.02\,\text{eV}$, one can see that for the special choices of ζ, see (109), only one of these dominates. For IO light neutrinos instead ϵ_2 behaves very similar to ϵ_1, because one of the two contributions clearly dominates. Regarding the width of the red and green areas in the different plots of figure 3 we see that these are mostly determined by the variation of the solar and atmospheric mass squared differences in their $3\,\sigma$ ranges, like in the case of the loop function $f(x)$, compare to figure 1. In the particular case of the CP asymmetry ϵ_2 for NO neutrino masses the reason for the broadness of the green band for larger values of m_0, see left plot in the middle of figure 3, is not this variation, but the fact that there are two different contributions to ϵ_2, see (84), that are of similar size in this parameter space. In the case of IO this does not happen, since the contribution accompanied by the loop function $f(m_1/m_2)$ is dominating over the other



Figure 4: **Case 1).** Baryon asymmetry Y_B of the Universe as a function of the lightest neutrino mass m_0 in the case of a light neutrino mass spectrum with NO (IO) in the left (right) panels. The plots at the top (bottom) are realized for $s = 1$ ($s = 3$) that predicts $\alpha = \pi/2$ ($3\pi/2$). This explains the different sign of Y_B for $s = 1$ than for $s = 3$. The other parameters are fixed as in figure 3. The horizontal blue band indicates the experimentally preferred $3\,\sigma$ range of the value of the baryon asymmetry $Y_B = (8.65 \pm 0.27) \times 10^{-11}$ [1]. The differently colored areas correspond to different choices of the parameter ζ shown in (109-111).

one. The sign of the CP asymmetries is determined by $\sin\alpha$ as well as the behavior of the loop function $f(x)$. For this reason, ϵ_1 can only be positive for small values of $m_0 \lesssim 4 \times 10^{-3}\,\text{eV}$, if light neutrinos follow NO, while ϵ_2 can only be positive for m_0 larger than 0.02 eV. In the case of ϵ_3 and NO the loop function instead does not change sign and thus ϵ_3 is always positive for this choice of s. Similarly, the CP asymmetries $\epsilon_{1,2}$ are always negative and can reach large values for light neutrinos with IO (indeed, the plots for ϵ_1 and ϵ_2 are almost identical in this case), as the relevant (dominating) loop function is always positive (negative) for $\epsilon_{1(2)}$. A suppression of $\epsilon_{1,2}$ occurs in this case for $\zeta \approx \pi/2,\ 3\pi/2$, see (83-85) and green areas in the upper- and middle-right panels of figure 3. In contrast, ϵ_3 is in this case positive for small $m_0 \lesssim 3 \times 10^{-2}\,\text{eV}$ and negative for larger values of m_0. It is, however, always strongly suppressed with respect to the other two CP asymmetries and thus irrelevant for the computation of Y_B (unless $\zeta \approx \pi/2,\ 3\pi/2$), since such a strong suppression cannot be compensated by the efficiency factor $\eta(m_i)$, see plot on the right of figure 2.

The described behavior of the CP asymmetries ϵ_i for NO and IO, respectively, explains the results obtained for the total baryon asymmetry Y_B, shown in the upper panels of figure 4. For NO, upper-left panel of figure 4, only values of $m_0 \approx 10^{-3}\,\text{eV}$ allow for the correct sign as well as size of Y_B for most of the values of ζ. Indeed, in about 80% of the parameter space $Y_B > 0$ is reproduced.[11] Moreover, Y_B within the experimentally preferred $3\,\sigma$ range, indicated by the blue

[11] We compute this percentage as follows: we take the interval of m_0 with $m_0 \lesssim 3 \times 10^{-3}\,\text{eV}$ and in which for some value of ζ the size of Y_B can be correctly achieved and calculate the size of the blue shaded area $I_{+(-)}$ with $Y_B > (<) 0$

band in figure 4, is achieved for $6.8 \times 10^{-4}\,\text{eV} \lesssim m_0 \lesssim 1.7 \times 10^{-3}\,\text{eV}$. The contribution Y_{B1}, arising from N_1 decays, dominates Y_B and is maximized for this particular range of m_0, since the efficiency factor $\eta(m_1)$ takes the largest value, see figure 2. For larger values of m_0, $m_0 \gtrsim 0.02\,\text{eV}$ instead Y_B can take positive values, but only for a rather small portion of the choices of ζ. In addition, its size is slightly too small compared to the experimentally preferred $3\,\sigma$ range for Y_B. This can be cured by choosing a value for $\tilde{\kappa}$ slightly larger than $\tilde{\kappa} = 4 \times 10^{-3}$, see (108). Since Y_B is proportional at LO to $\tilde{\kappa}^2$, an increase of $\tilde{\kappa}$ by less than a factor of two is sufficient. The situation strongly differs for the case of a light neutrino mass spectrum with IO, since there Y_B is mostly driven by the CP asymmetries ϵ_1 and ϵ_2 that are both negative for $s = 1$, see panels on the right of figure 3. Only for $m_0 \approx 3 \times 10^{-3}\,\text{eV}$ also positive values can be achieved. However, this occurs only in a small portion of the parameter space for ζ (less than 10%). Moreover, the size of Y_B is below the experimental lower $3\,\sigma$ limit (the maximum value achieved is about 6.5×10^{-11}). Thus, we conclude that for $s = 1$ and $k_1 = k_2 = 0$ only for NO light neutrino masses and $m_0 \approx 10^{-3}\,\text{eV}$ we can consider the sign as well as the size of Y_B to be correctly explained.

The situation is reverse in the case $s = 3$, since in this case the sign of $\sin\alpha$ is opposite. This is expected, since $s = 1$ is related to the choice $s = 3$ via the transformation where s is replaced by $n - s$, which changes the sign of all CP asymmetries, see (87). Thus, for $m_0 \gtrsim 3 \times 10^{-3}\,\text{eV}$ and NO of the light neutrino masses the sign (and also the size) of Y_B can be correctly achieved for most of the choices of ζ. Even more pronounced is the situation for IO, because for $s = 3$ the sign of Y_B is for (almost) all values of ζ correctly accommodated in the whole range of m_0 displayed, $5 \times 10^{-4}\,\text{eV} \lesssim m_0 \lesssim 0.1\,\text{eV}$. Also the size of Y_B can be correctly reproduced for some choice of ζ in the whole range of m_0. In figure 5, we also consider different choices of the parameters k_1 and k_2. In particular, we show in the plots on the left the effect of $k_2 = 1$. Most importantly, we see that for small values of m_0, $m_0 \lesssim 2.8 \times 10^{-3}\,\text{eV}$ and NO light neutrino masses Y_B is positive for all admitted values of ζ, see (109-110). This leads to a unique prediction of its sign compared to the case in which $k_2 = 0$ and thus the sign is correctly determined for all values of ζ. It can be traced back to the change in the CP asymmetry ϵ_2, compare left panel in the middle of figure 3 with the lower plot on the left of figure 5, while ϵ_1 does not change and ϵ_3, which is subdominant, changes sign because of $\eta(m_3)$ being much smaller than $\eta(m_{1,2})$. At the same time, Y_B is no longer positive for $m_0 \gtrsim 0.014\,\text{eV}$. On the right of figure 5 we see that for $k_1 = k_2 = 1$ and IO light neutrino masses Y_B behaves in the same way for $s = 1$ as for $s = 3$ and $k_1 = k_2 = 0$, see figure 4, showing clearly the importance of the choice of $k_{1,2}$. In particular, the choice $k_1 = 1$ is needed for achieving the correct sign of the baryon asymmetry. This value of k_1 and $s = 1$ entail $\alpha = 3\pi/2$ which is also obtained for $k_1 = 0$ and $s = 3$.

The approximate formulae given in subsection 3.3 describe in most of the parameter space the results obtained here with the formulae not expanded in $\tilde{\kappa}$ quite well. To quantify this statement better we first derive the next-to-leading order (NLO) terms in $\tilde{\kappa}$ contributing to the three CP asymmetries ϵ_i. The latter, expanded up to NLO in $\tilde{\kappa}$, can be written in the following compact

for this interval of m_0. The ratio $I_+/(I_+ + I_-)$ then corresponds to the percentage of parameter space in which Y_B is positive – in this case about 80%. We do so, since by changing the size of $\tilde{\kappa}$, we can in principle for all choices of ζ that lead to $Y_B > 0$ also achieve the correct size of Y_B, see (1). We use this measure also in the other cases to estimate the predictability of our approach regarding the sign of Y_B.



Figure 5: **Case 1).** Upper panels: Y_B versus m_0 for NO (left) and IO (right) spectrum. Lower panel: CP asymmetry ϵ_2 versus $m_0 = m_1$ in the case of NO. Group theoretical parameters: $n = 4$, $s = 1$ and different choices of k_1 and k_2. All the other parameters are the same as in figure 4. The red, green and light-blue areas correspond to the values of ζ reported in (109-111).

form[12]

$$\epsilon_1^{\rm NLO} = \epsilon_1^{\rm LO}\left[1+\frac{\tilde{\kappa}}{\sqrt{3}\,y_0}\sec(\theta+\zeta)\left(-\cos 3\,\theta+\frac{w^2}{z^2}\cos(\theta-2\,\psi)\right)\right], \tag{112}$$

$$\begin{aligned}\epsilon_2^{\rm NLO} = {} & -\epsilon_1^{\rm LO}\left[1+\frac{\tilde{\kappa}}{\sqrt{3}\,y_0}\sec(\theta+\zeta)\left(\cos(2\,\zeta-\theta)+\frac{w^2}{z^2}\cos(\theta-2\,\psi)\right)\right]\frac{f\,(m_2/m_1)}{f\,(m_1/m_2)} \\ & -\epsilon_3^{\rm LO}\left[1+\frac{\tilde{\kappa}}{\sqrt{3}\,y_0}\csc(\theta+\zeta)\left(-\sin(2\,\zeta-\theta)+\frac{w^2}{z^2}\sin(\theta-2\,\psi)\right)\right]\frac{f\,(m_2/m_3)}{f\,(m_3/m_2)},\end{aligned} \tag{113}$$

$$\epsilon_3^{\rm NLO} = \epsilon_3^{\rm LO}\left[1+\frac{\tilde{\kappa}}{\sqrt{3}\,y_0}\csc(\theta+\zeta)\left(\sin 3\,\theta+\frac{w^2}{z^2}\sin(\theta-2\,\psi)\right)\right] \tag{114}$$

where we have defined

$$\mathrm{Im}(z_1) = w\,\cos\psi \quad \text{and} \quad \mathrm{Im}(z_2) = w\,\sin\psi \tag{115}$$

with $w > 0$ being of order one and $0 \leq \psi < 2\,\pi$, since also the imaginary parts of the two complex parameters $z_{1,2}$ enter at this level. We have, furthermore, made use of the LO expressions $\epsilon_i^{\rm LO}$ of the CP asymmetries that can be found in (83-85). As one can see, the NLO terms are suppressed by $\tilde{\kappa}/y_0$ relative to the LO term. Leaving aside for the moment the effect of the imaginary parts of $z_{1,2}$, by setting $w = 0$, we can check that the NLO corrections are small compared to the LO term, if

$$\tilde{\kappa} \ll 0.07 \lesssim \sqrt{3}\,y_0 \tag{116}$$

assuming as allowed range of y_0 the one given in (69). With our choice of $\tilde{\kappa}$ in (108) this bound is clearly fulfilled. Consequently, we find in our numerical analysis that for all admitted values of ζ

[12]The appearance of $\sec(\theta+\zeta)$ and $\csc(\theta+\zeta)$ does not indicate any poles in $\epsilon_i^{\rm NLO}$, but arises only because we have extracted $\epsilon_i^{\rm LO}$. Indeed, the latter contain appropriate factors of $\cos(\theta+\zeta)$ and $\sin(\theta+\zeta)$ to cancel the former factors.

the relative difference between the LO approximations in (83-85) and the results obtained with the formulae not expanded in $\tilde{\kappa}$ is less than 10%. If we consider the impact of the imaginary parts of $z_{1,2}$, we see that for $w/z \lesssim 1$ the LO approximations still describe the unexpanded result very well. Indeed, in more than 90% of the parameter space of ζ, ψ and w with $w/z \lesssim 1$ the relative difference between the two is less than 10%. Only if w is taken to be larger than z, we find a considerable decrease in the goodness of the LO approximations, i.e. a relative deviation of the latter from the unexpanded results less than 10% is only given in about $60\% \div 70\%$ of the parameter space. Similar results are also obtained in the numerical analyses of case 2) and case 3 b.1) [as well as case 3 a)]. This shows that the simple LO expressions are quite powerful in describing in a correct way the CP asymmetries ϵ_i.

Leptogenesis in case 2)

As an explanatory example of case 2), we perform a numerical calculation of Y_B for the group theoretical parameters

$$n = 10 \quad \text{and} \quad u = 4\,. \tag{117}$$

It is important to note that in this case the matrix U_R (and thus also the PMNS mixing matrix) is given by the first matrix in (42), i.e. in order to achieve compatibility with the measured values of the mixing angles we have to apply a cyclic permutation of the rows of the original matrix in (36). As a consequence, when using the formulae for the mixing angles, found in [27], the CP invariants given in (39-40) and the LO approximations for the CP asymmetries ϵ_i in (89-91), we have to apply the set of transformations in (44). Clearly, if we had chosen to discuss a numerical example in which the correct form of U_R (and thus U_{PMNS}) is the second matrix in (42), we would need to apply the set of transformations in (45).

For completeness, we report the results of the lepton mixing angles, the phases α, β and δ, using $1.40 \lesssim \theta \lesssim 1.44$, in table 2. The latter interval of θ is also used in our numerical analysis. As discussed in [27], also $1.70 \lesssim \theta \lesssim 1.74$ is an admitted interval for θ, since the mixing angles can be accommodated equally well as for θ belonging to the other interval. However, $\sin\beta$ and $\sin\delta$ change sign, while $\sin\alpha$ remains invariant, and thus certain changes are observed in the results of the CP asymmetries ϵ_i.

The example of case 2) has a much richer phenomenology than the one studied for case 1), since all three CP phases are in general non-vanishing and, in addition, the Majorana phase α assumes three considerably different values for different choices of the group theoretical parameter v. As shown in table 2, the Dirac phase δ as well as the Majorana phase β are predicted to be almost maximal in this case, while the other Majorana phase α can take three different values (small, almost maximal, intermediate), depending on the choice of the group theoretical parameter v which can take five different values

$$v = 0,\, 6,\, 12,\, 18,\, 24 \tag{118}$$

for $u = 4$ and $n = 10$, obeying the constraint in and around (38). As one can read off from table 2, $v = 0$ leads to a small, $v = 6$, 24 to an almost maximal and $v = 12$, 18 to an intermediate value of $\sin\alpha$. Let us mention again that the sign of $\sin\alpha$ depends also here on the choice of the parameter k_1 and all values displayed in table 2 refer to $k_1 = 0$. In the same vein, the sign of $\sin\beta$ given in this table holds for $k_2 = 0$ and changes for $k_2 = 1$.

In figure 6 the variety of results for Y_B, its sign and its size, are shown for three different choices of the group theoretical parameter v. As mentioned, in case 2) we only display plots for a light neutrino mass spectrum with NO and always set $k_1 = k_2 = 0$. The smallness of $\sin\alpha$ for $v = 0$ entails a strong suppression of the otherwise dominating contribution to the CP asymmetries. A consequence of this suppression is that the sign of Y_B cannot be predicted in this case, but depends crucially on ζ, see (92) with ϕ_u being appropriately replaced according to (44). In this case, as can be checked by explicit computation, the Dirac phase δ provides the dominant source of CP violation in leptogenesis, yielding $|Y_B| \gtrsim 5 \times 10^{-11}$ for $m_0 \gtrsim 0.02$ eV (see the light-blue area in the upper-left panel of figure 6). As in the numerical example of case 1) discussed before, the red and

n	10		
u	4		
θ	$1.40 \div 1.44$		
$\sin^2\theta_{12}$	$0.340 \div 0.342$		
$\sin^2\theta_{13}$	$0.0187 \div 0.0250$		
$\sin^2\theta_{23}$	$0.558 \div 0.559$		
$\sin\beta$	$-0.94 \div -0.83$		
$\sin\delta$	$0.80 \div 0.86$		
v	0	6 , 24	12 , 18
$\sin\alpha$	$0.028 \div 0.035$	$0.94 \div 0.96$	$-0.62 \div -0.56$

Table 2: **Case 2)**. Results for lepton mixing angles and CP phases α, β and δ for $n = 10$, $u = 4$. All are compatible at the $3\,\sigma$ level or better with the experimental data. The values of $\sin\alpha$, $\sin\beta$ and $\sin\delta$ refer to the choice $k_1 = k_2 = 0$. The parameter v takes five different values: $v = 0$, 6, 12, 18 and 24. The interval of $\sin\alpha$ corresponding to $v = 18$ ($v = 24$) is the same as for $v = 12$ ($v = 6$). An atmospheric mixing angle in the first octant, $0.441 \lesssim \sin\theta_{23} \lesssim 0.442$, is obtained by a permutation of the second and third rows of the PMNS mixing matrix [27]. In this case $\sin\delta$ changes sign, while $\sin\alpha$ and $\sin\beta$ are invariant. For values of the parameter θ in the second admitted interval, $1.70 \lesssim \theta \lesssim 1.74$, compare [27], the mixing angles and $\sin\alpha$ are the same, but $\sin\beta$ and $\sin\delta$ change sign.

green bands represent the special choices $\zeta = 0$, π and $\zeta = \pi/2$, $3\pi/2$, respectively. These confirm our analytic findings, see (92), that a shift in ζ by $\pi/2$ changes the sign of all the CP asymmetries and therefore of Y_B. Notice that the red and green bands do not need to overlap with the light-blue area, since the intervals of ζ in (110) used to obtain the latter area do not contain the special values $\zeta = 0$, $\pi/2$, π, $3\pi/2$. Indeed, for values of $m_0 \lesssim 3 \times 10^{-3}\,\text{eV}$ not arbitrarily small values of Y_B can be achieved. The minimum possible values of $|Y_B|$ are obtained for $\zeta \approx 0.75$ and $\zeta \approx 2.32$. The vanishing of Y_B for $m_0 \approx 3 \times 10^{-3}\,\text{eV}$ is due to the vanishing of the loop function $f(m_1/m_2)$, see upper-left panel of figure 1. Although for large enough m_0, $m_0 \gtrsim 2.7 \times 10^{-2}\,\text{eV}$ Y_B sufficiently large can be obtained, we thus do not consider this case as a successful one.

In contrast, for $v = 6$ the sign of Y_B can be univocally be predicted for $m_0 \lesssim 3 \times 10^{-3}\,\text{eV}$ and is in accordance with the experimental observations. The dominant contribution to the baryon asymmetry (with the correct sign) for NO is produced by the out-of-equilibrium decays of the lightest RH neutrino, N_1, as expected from the general results given in subsection 3.2. For our particular choice of $\tilde{\kappa}$ in (108) we see that in the interval $7 \times 10^{-4} \lesssim m_0 \lesssim 2 \times 10^{-3}$ the size of Y_B can be correctly achieved. We also see that for $m_0 \gtrsim 3 \times 10^{-3}\,\text{eV}$ the sign of Y_B is almost always negative. If we consider another choice of the parameters $k_{1,2}$ as in (105), we can make Y_B positive for $m_0 \gtrsim 3 \times 10^{-3}\,\text{eV}$. The situation is reverse for the choice $v = 12$, simply because the sign of $\sin\alpha$ is opposite compared to the one in the case $v = 6$. So, for $v = 12$ the baryon asymmetry Y_B is always negative for $m_0 \lesssim 2 \times 10^{-3}\,\text{eV}$ and almost always positive for $m_0 \gtrsim 2 \times 10^{-3}\,\text{eV}$. Consequently, only for values of m_0 larger than $6 \times 10^{-3}\,\text{eV}$ also the correct size of Y_B can be obtained for a certain choice of ζ, if we keep $\tilde{\kappa}$ fixed to the value as in (108). In the range $6 \times 10^{-3}\,\text{eV} \lesssim m_0 \lesssim 0.1\,\text{eV}$ positive Y_B is obtained for more than 90% of the choices of ζ. Again, like in the case $v = 6$ the sign of Y_B can be changed for also $v = 12$ by changing the values of the parameters $k_{1,2}$. Comparing the results for $v = 6$ and $v = 12$ we note in addition that the size of $|Y_B|$ is a bit smaller for $v = 12$ than for $v = 6$, simply because $|\sin\alpha|$ is smaller for $v = 12$ than for $v = 6$, see table 2.

Concerning the IO spectrum, the results are very similar to the ones discussed in case 1), that is the dominant contribution to the baryon asymmetry comes from N_1 and N_2 decays, which can generate large CP asymmetries of order few times 10^{-5} and with the correct sign. For this reason we do not report the corresponding plots of Y_B. We only remark that, as for the NO spectrum, in



Figure 6: **Case 2).** Y_B versus m_0 for $n = 10$ and $u = 4$ and three choices of v. Results for $v = 18$ ($v = 24$) are equal to the plot with $v = 12$ ($v = 6$). In all the plots $1.40 \lesssim \theta \lesssim 1.44$ and the neutrino mass squared differences are varied within their experimentally preferred $3\,\sigma$ ranges. The horizontal blue band represents the measured value of Y_B at the $3\,\sigma$ level. The red, green and light-blue areas correspond to the choices of ζ reported in (109-111).

the case $v = 0$, the sign of Y_B is not fixed, see (92), while for other choices of v the sign can be determined.

Leptogenesis in case 3)

A representative example for case 3) is given by

$$n = 8 \quad \text{and} \quad m = 4 \tag{119}$$

which leads to good agreement with the experimental data on lepton mixing angles, if we consider case 3 b.1), as has been shown in [27]. The discussion of this case has two advantages: the value of the index n of the group $\Delta(6\,n^2)$ is still moderately small and thus the group itself is not too large (it has 384 elements) and, at the same time, several values of the group theoretical parameter s, characterizing the CP transformation X, are admitted by the requirement to accommodate all three lepton mixing angles at the $3\,\sigma$ level or better, so that different types of results for leptogenesis are achieved. Concretely, the viable choices of s are [27]

$$s = 1,\, 2,\, 4 \tag{120}$$

as well as the values of s with $s > n/2 = 4$ that are related to the mentioned ones via $n - s = 8 - s$, i.e. $s = 6$ and $s = 7$. These are not discussed independently, since results for these cases can be deduced from those for $s \leq n/2 = 4$, see table 6 and explanations in [27] regarding the mixing parameters and (98) for the CP asymmetries ϵ_i. The results of mixing angles and CP phases for the different choices of s are summarized in table 3. As one can see, all values of s give rise to a similar fit to the solar and the reactor mixing angles, while the results obtained for θ_{23} differ. The value of the latter also depends on the interval chosen of the parameter θ, since replacing θ by $\pi - \theta$ changes

n	8		
m	4		
$\sin^2\theta_{12}$	$0.316 \div 0.321$		
$\sin^2\theta_{13}$	$0.0186 \div 0.0250$		
s	1	2	4
θ	$1.29 \div 1.33$	$1.81 \div 1.82$	$1.29 \div 1.33$
$\sin^2\theta_{23}$	$0.416 \div 0.427$	$0.635 \div 0.643$	$1/2$
$\sin\alpha = \sin\beta$	$-1/\sqrt{2}$	1	0
$\sin\delta$	$0.934 \div 0.937$	$0.734 \div 0.738$	-1

Table 3: **Case 3 b.1)**. Lepton mixing angles and CP phases α, β and δ for the choice $n = 8$ and $m = 4$. The group theoretical parameter s, characterizing the CP transformation X, can take different values. Here we display those leading to good agreement (at the $3\,\sigma$ level or better) with the experimental results on the lepton mixing angles and that fulfill $s \leq n/2 = 4$. Results for $s > n/2 = 4$ can be easily deduced from the ones shown in this table using the transformation in (54). The equality of the absolute values of $\sin\alpha$ and $\sin\beta$ originates from $m = n/2 = 4$, see (51). Setting $k_1 = k_2 = 0$ leads to $\sin\alpha = \sin\beta$. Using (51) we can also check that $s = n/2 = 4$ entails trivial Majorana phases and a maximal Dirac phase. We display only one interval for θ. However, in most cases (probably for $s = 2$ not) θ can be in a second admitted interval. This is related to the one shown via the transformation of θ in $\pi - \theta$. In this case, $\sin^2\theta_{23}$ becomes $\cos^2\theta_{23}$ and $\sin\delta$ changes sign. For further details see [27].

$\sin^2\theta_{23}$ into $\cos^2\theta_{23}$. Consequently, for most of the choices of s two intervals of θ are admitted that lead to different results not only for the atmospheric mixing angle, but also to a different sign in $\sin\delta$. In our numerical analysis of leptogenesis, we will always stick to choices of θ belonging to the intervals displayed in table 3 and only briefly comment on results originating from other choices of θ. A feature of this example is $m = n/2 = 4$ which leads to a considerable simplification of the formulae for the CP phases, see (51). From there we see immediately that $\sin\alpha$ and $\sin\beta$ must be equal up to a sign (for the particular choice $k_1 = k_2 = 0$ they are equal) for all choices of s and, moreover, that for $s = n/2 = 4$ both Majorana phases are trivial, whereas the Dirac phase is maximal. As shown in table 3 in our particular example the Dirac phase is large in all cases and also the Majorana phases are sizable, if they are not trivial. Furthermore, we note that $\sin\alpha$ and $\sin\beta$ turn out to be negative for $s = 1$, while they are positive for $s = 2$. All these observations are relevant for the prediction of the sign of the baryon asymmetry Y_B.

We report the results for the baryon asymmetry Y_B depending on the light neutrino mass m_0 and for the different choices $s = 1$, $s = 2$ and $s = 4$ in figure 7. Like for case 2), we focus on a light neutrino mass spectrum with NO. Again, the areas in different colors stand for different choices of the parameter ζ, see (109-111). As can be seen, the correct sign and size can be achieved for all three values of s. For $s = 1$ the sign of Y_B is mostly negative for small m_0, $m_0 \lesssim 3 \times 10^{-3}\,\text{eV}$. In fact, for such values of the lightest neutrino mass we have $\epsilon_1 < 0$ and $\epsilon_2 > 0$. Then, the dominant contribution to the baryon asymmetry (with a negative sign) comes from the heaviest RH neutrino, N_1, provided $|\epsilon_1|$ is not strongly suppressed (this suppression happens for $\zeta \approx \pi/2,\ 3\pi/2$). In the interval 3×10^{-3} eV $\lesssim m_0 \lesssim 10^{-2}$ eV both ϵ_1 and ϵ_2 are positive and thus the overall sign of Y_B. This is also true for $m_0 \gtrsim 10^{-2}$ eV as long as the contribution coming from N_1 is not suppressed due to the particular choice of ζ. Thus, we correctly predict the sign of Y_B in more than 90% of the parameter space for 3×10^{-3} eV $\lesssim m_0 \lesssim 0.1$ eV. Regarding the size of Y_B, this is correctly achieved for $\tilde{\kappa} = 4 \times 10^{-3}$, as long as $m_0 \gtrsim 5 \times 10^{-3}$ eV. The results for $s = 2$ are similar to those for $s = 1$, with the sign of Y_B reversed, because the dominant contribution to the CP asymmetries that is proportional to $\sin\alpha$ changes sign, as $\sin\alpha$ is negative for $s = 1$ and positive for $s = 2$, see table 3. Consequently, only for small values of m_0 the sign is correctly

reproduced. Y_B compatible with its measured value at the $3\,\sigma$ level or better prefers m_0 in the interval 6.8×10^{-4} eV $\lesssim m_0 \lesssim 1.7 \times 10^{-3}$ eV. In this particular interval, the sign of Y_B is positive in more than 80% of the parameter space. The size of the maximally achievable value of $|Y_B|$ is slightly smaller for $s = 1$ than for $s = 2$, simply because the dominant term is proportional to $\sin\alpha$ that is smaller by a factor $1/\sqrt{2} \approx 0.71$ for $s = 1$ than for $s = 2$, compare also table 3. The remaining choice $s = 4$ allows to achieve the correct sign and size of Y_B in some parameter space for larger m_0, for which light neutrino masses become more and more degenerate. However, like for $v = 0$ in case 2), we cannot call this a prediction of the sign of Y_B, since there is no preference for a certain sign observable in figure 7. This is explained by the absence of the leading term(s) in the CP asymmetries that are sourced by non-trivial Majorana phases. A way to see this clearly is to use the formulae in (99-101) that are derived under the assumption that $m = n/2$ and $s = n/2$ in case 3 b.1). As one can see, the resulting CP asymmetries are proportional to $\sin 2\zeta$ as well as to $\sin 2\theta$. Thus, considering ζ in the intervals in (110) leads to equally likely to $\sin 2\zeta > 0$ than to $\sin 2\zeta < 0$. Consequently, the sign of Y_B cannot be predicted in this case, only by fixing the group theoretical parameters, k_1, k_2 and the interval of θ. In addition, we note that also a change in the interval of θ gives rise to a change in the sign of the CP asymmetries, since both intervals are connected by replacing θ with $\pi - \theta$. According to the approximations in (99-101) the CP asymmetries are very small for the special choices $\zeta = 0$, π and $\zeta = \pi/2$, $3\pi/2$, represented by the red and green bands in the figures. As already mentioned in subsection 3.3, for the first choice the CP asymmetries vanish exactly to all orders in $\tilde{\kappa}$, whereas for the second choice this statement is only true at LO. Nevertheless, also in the latter case the CP asymmetries become way too small for explaining the experimentally observed value of Y_B. The light-blue area is delimited by the curves defined by $\zeta = \pi/4$ ($\zeta = 3\pi/4$) and $\zeta \approx 0.24$ ($\zeta \approx 1.82$), respectively. The former two choices refer to the maximally achievable values of Y_B (with positive and negative sign, respectively). This can be easily understood by using the formulae in (99-101) which show that the LO expressions of all CP asymmetries ϵ_i are maximized for maximal $|\sin 2\zeta|$ which occurs, for example, for $\zeta = \pi/4$ and $\zeta = 3\pi/4$. The latter two choices describe the boundaries of the light-blue areas that lead to the smallest absolute values for Y_B for a certain value of m_0. They simply correspond to two borders of the intervals $\mathcal{I}_i$ in which ζ can vary, if it does not take special values, compare (110).

It is interesting to compare in more detail the results for $s = 4$ in this case with the ones obtained for $v = 0$ in case 2), see figure 6. As said, they share the feature that the sine of the Majorana phase α is small (or even exactly zero), see table 2 and 3, and thus the otherwise dominant contribution to the CP asymmetries ϵ_i is absent. Furthermore, the Dirac phase is in both cases large, $|\sin\delta| \gtrsim 0.8$. However, there is also a crucial difference between them, namely the fact for $v = 0$ in case 2) the Majorana phase β is large, whereas it is trivial for $s = 4$ in this case. Thus, comparing these two cases allows us to understand the effect of the CP phases β and δ separately on the resulting baryon asymmetry. Indeed, we see that the two figures, upper-left plot of figure 6 and lower-left plot of figure 7, are quite similar, with the striking difference that for $m_0 \lesssim 2 \times 10^{-3}$ eV for $s = 4$, $\sin\beta = 0$, Y_B is very small, while Y_B can reach values up to $\pm 0.3 \times 10^{-11}$ for $v = 0$ in case 2), i.e. $0.83 \lesssim |\sin\beta| \lesssim 0.94$.

Finally, we comment on some peculiarities of examples belonging to case 3.a): clearly, for (the always admitted) choice $s = 0$ no non-vanishing CP asymmetry can be obtained, because in this case an accidental CP symmetry, common to charged lepton and neutrino sectors, exists, for details see [27]. Furthermore, we note that in some cases such an accidental CP symmetry arises from a particular choice of the parameter θ, see e.g. $n = 16$, $m = 1$ and $s = 8$ in which the best fitting point of θ is at $\theta_{\rm bf} = 0$. Considering the whole $3\,\sigma$ ranges of the mixing angles also non-zero values of θ are compatible with the experimental data. However, we expect then that CP phases are still small and thus the size of Y_B is suppressed. Eventually, we notice that there are cases in which $\sin\alpha = 0$ is obtained at the best fitting point $\theta_{\rm bf}$ (indeed, in these cases the best fit value of the solar mixing angle cannot be accommodated). These are cases in which a prediction of the sign of Y_B is impossible, since we expect a similar behavior as for $s = 4$ for case 3 b.1), as discussed here, or for $v = 0$ in case 2).



Figure 7: **Case 3 b.1).** Y_B versus m_0 for $n = 8$ and $m = 4$. We display results for $s = 1$, 2, 4. Those for $s > n/2 = 4$ can be deduced from the ones presented here. The parameter θ lies in the intervals given in table 3. Differently colored areas indicate different choices of ζ, see (109-111). The horizontal blue band corresponds to the experimentally preferred $3\,\sigma$ range of Y_B [1]: $Y_B = (8.65 \pm 0.27) \times 10^{-11}$.

4 Comments on flavored leptogenesis

In this section we briefly comment on the case of flavored leptogenesis which is realized if RH neutrino masses are smaller than 10^{12} GeV [50–52] (and larger than 10^8 GeV in order to correctly reproduce the light neutrino mass scale [53] for $y_0 \gtrsim 10^{-3}$). In particular, for $10^9\,\text{GeV} \lesssim M_i \lesssim 10^{12}\,\text{GeV}$ the Yukawa interactions of RH charged leptons of τ-flavor are in thermal equilibrium and hence leptogenesis occurs in the so-called two-flavor regime. For even lower RH neutrino masses, $M_i \lesssim 10^9$ GeV, also the μ-flavor becomes dynamical.

The formula for the flavored CP asymmetry ϵ_i^α ($\alpha = e,\, \mu,\, \tau$) reads [49]

$$\begin{aligned}
\epsilon_i^\alpha &= -\frac{\Gamma(N_i \to H l_\alpha) - \Gamma(N_i \to H^\star \bar{l}_\alpha)}{\sum_\beta [\Gamma(N_i \to H l_\beta) + \Gamma(N_i \to H^\star \bar{l}_\beta)]} \\
&= -\frac{1}{8\pi\,(\hat{Y}_D^\dagger \hat{Y}_D)_{ii}} \sum_{j \neq i} \left\{ \text{Im}\left((\hat{Y}_D^\dagger \hat{Y}_D)_{ij} (\hat{Y}_D)^*_{\alpha i} (\hat{Y}_D)_{\alpha j} \right) f(M_j/M_i) \right. \\
&\quad \left. + \text{Im}\left((\hat{Y}_D^\dagger \hat{Y}_D)_{ji} (\hat{Y}_D)^*_{\alpha i} (\hat{Y}_D)_{\alpha j} \right) g\,(M_j/M_i) \right\}, \qquad (121)
\end{aligned}$$

where $f(x)$ in the SM is given in (59) and

$$g(x) \;=\; \frac{1}{1 - x^2}\,. \qquad (122)$$

Notice that the term in (121) with $g(x)$ does not depend on Majorana phases, because it originates from the lepton number conserving, but lepton flavor violating one-loop contribution to the CP asymmetries. The analytic expression of the unflavored CP asymmetry ϵ_i in (58) is recovered, if we sum over the flavor index α

$$\epsilon_i \;=\; \sum_{\alpha = e,\,\mu,\,\tau} \epsilon_i^\alpha\,. \qquad (123)$$

The contribution to Y_B from the decay of each RH neutrino has a form similar to expression (57), but with ϵ_i^α instead of ϵ_i and efficiency factors with an additional index, depending on the dynamical flavor.

In analogy to the study of leptogenesis performed in the unflavored regime, we consider a neutrino Yukawa matrix Y_D of the form $Y_D = y_0\,\mathbb{1} + \delta Y_D$, with the correction δY_D defined as in (27) and proportional to the small expansion parameter κ ($\tilde{\kappa}$). We can first determine the allowed interval of the parameter y_0 using (16) and taking into account 10^9 GeV $\lesssim M_i \lesssim 10^{12}$ GeV

$$0.001 \lesssim y_0 \lesssim 0.04\,. \tag{124}$$

Applying (121) we can check that for our form of Y_D the flavored CP asymmetries ϵ_i^α are proportional to the product of y_0 and κ ($\tilde{\kappa}$) at LO. Thus, corrections δY_D are also in this case crucial in order to achieve non-vanishing CP asymmetries. However, the dependence on the parameters y_0 and κ is different than in the unflavored case in which ϵ_i are independent of y_0 and scale as κ^2. Since (123) holds, we see that the sum of ϵ_i^α over α must add up to zero at LO. Hence, we can always express one of the flavored CP asymmetries as the negative of the sum of the other two at this order. For y_0 varying within the interval in (124) the absolute value of the CP asymmetries ϵ_i^α is larger than 10^{-6}, if we choose $\tilde{\kappa} \gtrsim 10^{-3}$. This lower bound coincides with the one necessary to obtain successful leptogenesis in the unflavored regime, see (79). As explained, this is also the expectation of the natural size of κ from model building. Another important feature of the LO result of the flavored CP asymmetries, common to the case of unflavored CP asymmetries, is the fact that the only source of CP violation relevant for leptogenesis is provided by the phases in the matrix U_R, coinciding with the phases of the PMNS mixing matrix. The reason for this to happen is the symmetric form of δY_D in flavor space so that the quantity $(\hat{Y}_D^\dagger \hat{Y}_D)_{ij}$, $i \neq j$, in (70) only depends on the real part of δY_D at LO.

We first derive the predictions for the flavored CP asymmetries, assuming that $U_R = U_{PMNS}$ with U_{PMNS} being of its general form as in (166). In order to simplify the resulting formulae we assume in the following $\zeta = k_\zeta\,\pi$ with $k_\zeta = 0,\,1$ (equivalent to vanishing real part of z_2) and the low energy CP violation to be encoded only in the Dirac phase δ, i.e.

$$\alpha \;=\; k_\alpha\,\pi \quad \text{and} \quad \beta \,+\, 2\,\delta \,=\, k_\beta\,\pi, \quad \text{with} \quad k_{\alpha,\beta} = 0,\,1\,. \tag{125}$$

At LO in $\tilde{\kappa}$, the flavored CP asymmetries ϵ_1^α of N_1 decays then read

$$\begin{aligned}
\epsilon_1^e &\approx (-1)^{k_\beta+k_\zeta}\,\frac{\sqrt{3}\,y_0\,\tilde{\kappa}}{16\,\pi}\,\cos^2\theta_{12}\,\sin^2 2\theta_{13}\,\sin 2\delta\, f\left(\frac{m_1}{m_3}\right)\,,\\
\epsilon_1^\mu &\approx (-1)^{k_\zeta+1}\,\frac{\sqrt{3}\,y_0\,\tilde{\kappa}}{4\,\pi}\,J_{CP}\,\left[(-1)^{k_\alpha}\,f\left(\frac{m_1}{m_2}\right)+(-1)^{k_\beta}\,f\left(\frac{m_1}{m_3}\right)+g\left(\frac{m_1}{m_2}\right)-g\left(\frac{m_1}{m_3}\right)\right]-\,\sin^2\theta_{23}\,\epsilon_1^e\,,\\
\epsilon_1^\tau &= -\epsilon_1^e\,-\,\epsilon_1^\mu\,,
\end{aligned} \tag{126}$$

where the CP invariant $J_{CP} \propto \sin\delta$ is defined in (168) and we have used relation (16). First of all, this example confirms that ϵ_i^α are proportional to the product $y_0\,\tilde{\kappa}$. It also reflects the property that the contribution containing $g(x)$ must be independent of Majorana phases, in particular of the low energy ones α and β, appearing in the PMNS mixing matrix. Furthermore, this example clearly shows that, unlike in the case of unflavored asymmetries, now the sign of the CP asymmetries (and Y_B) cannot be predicted just from the knowledge of the sign of the sine of the CP phases, because it explicitly depends on k_ζ which encodes the sign of the real part of the correction z_1. Finally, we note that in this particular case $\epsilon_1^e \approx 0$ and $\epsilon_1^\tau \approx -\epsilon_1^\mu \propto J_{CP}$, if the CP phase δ is (close to) maximal, $\delta \approx \pi/2,\,3\pi/2$. The form of ϵ_2^α and ϵ_3^α is very similar to the one of ϵ_1^α, in particular, regarding relations between the CP asymmetries of the different flavors α as well as the dependence on the CP phase δ.

Concerning the different scenarios with a flavor and a CP symmetry we focus on the flavored

CP asymmetries for case 1), since simple expressions can be derived. We obtain for $\epsilon^\alpha_{1,3}$ at LO

$$\begin{aligned}
\epsilon^\alpha_1 &\approx (-1)^{k_1+1}\,\frac{y_0\,\tilde{\kappa}}{6\sqrt{3}\,\pi}\,\cos(\theta+\zeta)\,\cos\left(\theta+\rho_\alpha\,\frac{\pi}{3}\right)\,\sin 6\,\phi_s\,f\left(\frac{m_1}{m_2}\right)\,,\\
\epsilon^\alpha_3 &\approx (-1)^{k_1+k_2+1}\,\frac{y_0\,\tilde{\kappa}}{6\sqrt{3}\,\pi}\,\sin(\theta+\zeta)\,\sin\left(\theta+\rho_\alpha\,\frac{\pi}{3}\right)\,\sin 6\,\phi_s\,f\left(\frac{m_3}{m_2}\right)\,,
\end{aligned} \tag{127}$$

with $\rho_e = 3$, $\rho_\mu = 1$ and $\rho_\tau = -1$. The formulae for ϵ^α_2 are given by the linear combination in (82), replacing $\epsilon_{1,3}$ with the flavored counterparts $\epsilon^\alpha_{1,3}$. Notice that the contribution to the CP asymmetries proportional to $g(x)$ is absent in this case. This is due to the fact that the Dirac phase is trivial, see (34). Most importantly, the sign of ϵ^α_i (and Y_B) depends on the particular choice of the parameter ζ (the ratio between the real parts of z_1 and z_2 that parametrize the correction δY_D, see (73) and (27)). This is in contrast to the unflavored case where ζ appears only as argument of positive semi-definite functions, cf. (83-85).

If Y_D is only invariant under the residual symmetry G_ν instead of the full flavor group G_f and the CP symmetry, ϵ^α_i are in general non-zero even for vanishing corrections δY_D. For Z and X representing the residual Z_2 and the CP symmetry in $\mathbf{3}$, Y_D is in this case constrained by the conditions

$$Z^\dagger\,Y_D\,Z \;=\; Y_D \quad \text{and} \quad X^\star\,Y_D\,X \;=\; Y_D^\star\,, \tag{128}$$

such that it can be in general written as

$$Y_D = \Omega\,R_{ij}(\theta_L)\,\begin{pmatrix} y_1 & 0 & 0\\ 0 & y_2 & 0\\ 0 & 0 & y_3\end{pmatrix}\,R_{ij}(-\theta_R)\,\Omega^\dagger \tag{129}$$

with $R_{ij}(\theta_{L,R})$ being rotations in the (ij)-plane around $\theta_{L,R}$, lying in the interval $[0,\pi)$, and y_i real parameters. All of these five are not further restricted by G_ν. As a consequence, the light neutrino masses as well as the lepton mixing angles and low energy CP phases are less correlated with the parameters appearing in M_R in (11). In particular, U_ν and thus U_{PMNS} depends on an undetermined angle $\tilde{\theta}$, different from θ and $\theta_{L,R}$. Furthermore, the relation between the light and heavy neutrino masses in (16) does not hold except for the light neutrino mass m_k with $k \neq i,j$. Only for this generation the CP parity of the light neutrino is given by the one of the RH one N_k, encoded in the matrix K_ν.

Transforming Y_D in (129) to the mass basis of RH neutrinos using U_R in (12), we find

$$\hat{Y}_D \;=\; \Omega\,R_{ij}(\theta_L)\,\begin{pmatrix} y_1 & 0 & 0\\ 0 & y_2 & 0\\ 0 & 0 & y_3\end{pmatrix}\,R_{ij}(\theta-\theta_R)\,K_\nu\,. \tag{130}$$

As can be checked, the flavored CP asymmetries, computed with (121), are in general proportional to $\sin 2(\theta-\theta_R)$ (but not θ_L), the product $y_i\,y_j$ as well as the difference $y_i^2 - y_j^2$. Thus, if any of these vanishes, i.e. $\theta = \theta_R + l\,\pi/2$, l integer, or $y_i = 0$ or $y_j = 0$ or $y_i = \pm y_j$, the CP asymmetries all must vanish. Furthermore, ϵ^α_k with $k \neq i,j$ is always zero. The correct size $|\epsilon^\alpha_i| \gtrsim 10^{-6}$ can be achieved for $y_j \gtrsim 10^{-3}$. This constraint on the neutrino Yukawa couplings is consistent with the requirement to reproduce light neutrino masses of order 0.1 eV. Interestingly enough, computing the unflavored CP asymmetries ϵ_i we see that they vanish exactly in this scenario, unless a small correction δY_D, which is not invariant under G_ν, is introduced.

In addition to this general discussion, we have explicitly studied the predictions for ϵ^α_i for all cases 1) to 3 b.1). In case 1) not only the unflavored CP asymmetries, but also the flavored ones vanish identically. For this reason, we present the exact expressions of ϵ^α_i for case 2)

$$\begin{aligned}
\epsilon^\alpha_{1\,(3)} &= \frac{y_1\,y_3\,(y^2_{1\,(3)} - y^2_{3\,(1)})\,\sin 2\,(\theta-\theta_R)\,\sin\left(\phi_u - \rho_\alpha\,\frac{\pi}{3}\right)}{48\,\pi\left(y^2_{1\,(3)}\,\cos^2(\theta-\theta_R) \,+\, y^2_{3\,(1)}\,\sin^2(\theta-\theta_R)\right)}\left[(-1)^{k_2}\,f\left(\frac{M_{3\,(1)}}{M_{1\,(3)}}\right) + g\left(\frac{M_{3\,(1)}}{M_{1\,(3)}}\right)\right]\,,\\
\epsilon^\alpha_2 &= 0\,.
\end{aligned} \tag{131}$$

with ρ_α defined as below (127). All characteristics mentioned above can be verified: $\epsilon_2^\alpha = 0$, since the rotation $R_{13}(\theta)$ acts in the (13)-plane; furthermore, $\epsilon_{1,3}^\alpha$ are proportional to the product $y_1\, y_3$, to the difference $y_1^2 - y_3^2$ as well as to $\sin 2\,(\theta - \theta_R)$; and the sum over all flavors α, see (123), vanishes. In addition, we observe that $\epsilon_{1,3}^\alpha$ are not sensitive to the parameter ϕ_v, which determines the Majorana phase α, see (41).[13] This result is in stark contrast with the ones of the unflavored CP asymmetries in (89-91) where the terms with ϕ_v usually dominate, if v does not vanish. Again, the sign of Y_B is not determined, since $y_{1,3}$ can be both positive and negative. This observation is contrary to the findings in the case of unflavored leptogenesis, see figure 6. The formulae for ϵ_i^α for case 3 a) and case 3 b.1) show a similar structure as those in (131) and are thus not explicitly discussed here.

Finally, we remark that for the particular choice $\phi_u = 0,\ \pi$ $(u = 0,\, n)$, the flavored CP asymmetries in case 2) in (131) fulfill

$$\epsilon^e_{1\,(3)} \;=\; 0 \quad \text{and} \quad \epsilon^\tau_{1\,(3)} \;=\; -\epsilon^\mu_{1\,(3)}\,. \tag{132}$$

This result is related to the presence of $\mu\tau$ reflection symmetry [22], since $X = P_{23}$ or $Z\,X = P_{23}$, if we set in addition $\phi_v = 0$ $(v = 0)$, compare (28). A similar result can also be obtained for case 3 b.1): if we set $m = n/2$, we find $\epsilon^e_{2\,(3)} = 0$ and $\epsilon^\mu_{2\,(3)} = -\epsilon^\tau_{2\,(3)}$ (the CP asymmetries ϵ_1^α vanish in all flavors due to the form of the rotation $R_{ij}(\theta)$). The $\mu\tau$ reflection symmetry is then recovered for $s = n/2$ [27] and we find explicitly $ZX = P_{23}$.[14]

Flavored leptogenesis in a scenario with $\mu\tau$ reflection symmetry has also been discussed in [54]. In their analysis the authors only impose this symmetry on the neutrino mass matrix. A diagonal mass matrix for charged leptons is ensured with gauged $L_\mu - L_\tau$ symmetry. The authors stick to the two-flavor regime and discuss the case in which the RH neutrino mass spectrum is strongly hierarchical. Thus, only the out-of-equilibrium decays of the lightest RH neutrino are relevant for generating the baryon asymmetry of the Universe.

5 Comments on SUSY framework

Here we comment on the implementation of our scenario and results for leptogenesis in a SUSY framework, since most of the concrete models with a flavor (and a CP) symmetry are formulated in a SUSY context, see e.g. [23, 41]. The relevant superpotential of such a scenario reads

$$w_l = Y_l\, l\, h_d\, \alpha^c + Y_D\, l\, h_u\, \nu^c + M_R\, \nu^c\, \nu^c \tag{133}$$

with $h_{u,d}$ denoting the two Higgs multiplets of the MSSM. The latter are taken to transform trivially under all non-gauge symmetries. We assign the three generations of RH neutrinos ν^c to $\mathbf{3}$ under G_f, while the fields l now transform as $\bar{\mathbf{3}}$. In this way, the term $l\, h_u\, \nu^c$ is invariant under G_f and thus arises at the renormalizable level. In addition, the Yukawa coupling Y_D is proportional to the identity matrix. The RH charged leptons α^c are in the trivial representation of G_f. Like in the non-SUSY framework, also here we assume the existence of an auxiliary symmetry $Z_3^{(\text{aux})}$. Only the two fields μ^c and τ^c carry non-trivial phases under the latter symmetry: $\mu^c \sim \omega^2$ and $\tau^c \sim \omega$.

We note a few crucial differences relevant for our results of leptogenesis. First of all, the range of RH neutrino masses in which unflavored thermal leptogenesis is the dominant generation mechanism of the lepton asymmetry is rescaled by the factor $1 + \tan^2\beta$ with $\tan\beta = \langle h_u\rangle/\langle h_d\rangle$ so that we have to require RH neutrino masses M_i to lie in the interval $10^{12}\,(1 + \tan^2\beta)\,\text{GeV} \lesssim M_i \lesssim 10^{14}\,(1 + \tan^2\beta)\,\text{GeV}$. We can estimate the allowed values of $\tan\beta$ in our scenario by considering the operator

[13] Although the form of Y_D is now more complicated than in the previous case and the PMNS mixing matrix does not coincide anymore with U_R, it is still true that the Majorana phase α depends in this case (almost) only on ϕ_v, as long as the measured lepton mixing angles are accommodated well.

[14] We note that this choice is also possible for case 3 a), since the latter arises from the same type of residual symmetries Z and X, see (28). However, it is clear that only for case 3 b.1) agreement with the experimental data on mixing angles can be achieved for the choice $m = n/2$ (and $s = n/2$) [27].

that gives rise to the tau lepton mass. This operator requires (at least) one insertion of a flavor symmetry breaking field, since LH leptons transform as $\overline{\mathbf{3}}$ under G_f, whereas τ^c and h_d are trivial singlets. Assuming that the size of the suppression due to the necessary insertion of one flavor symmetry breaking field is $\varepsilon \approx (0.01 \div 0.1)$, see (192) in appendix D, and that the tau Yukawa coupling varies between 1/3 and 3, we get as range of $\tan\beta$ the following[15]

$$\begin{aligned} \varepsilon = 0.01 \quad &: \quad \tan\beta \approx 3 \,, \\ \varepsilon = 0.05 \quad &: \quad 3 \lesssim \tan\beta \lesssim 15 \,, \\ \varepsilon = 0.1 \quad &: \quad 3 \lesssim \tan\beta \lesssim 30 \,. \end{aligned}$$

Thus, RH neutrino masses are expected to be larger than in the SM case. In order to obtain the correct values for the light neutrino masses we have to re-scale the coupling y_0 accordingly, see (16).

On the other hand, the masses of RH neutrinos cannot be too large in a SUSY framework, since larger values give rise to larger contributions to flavor non-universal soft terms of sleptons through renormalization group running effects [55]. These flavor non-universal soft terms play a crucial role in charged lepton flavor violating processes [56] such as $\mu \to e\gamma$ and $\mu - e$ conversion that are strongly constrained experimentally [57, 58] (for a review see [59]).

Furthermore, large RH neutrino masses also require a large reheating temperature $T_R \gtrsim 10^{12}$ GeV. This in turn gives rise to the well-known gravitino problem [60], unless e.g. the gravitino production is suppressed and/or it decays [61] and/or there is an additional substantial contribution to the total energy density of the Universe that dilutes the gravitino abundance, see discussion in [46].

In a SUSY framework the CP asymmetries $\epsilon_i^{(\alpha)}$ not only arise from decays of the RH neutrinos ν_i^c to SM particles, but also from decays of their SUSY partners and to SUSY particles, see e.g. [46]. Thus, also the form of the loop functions $f(x)$ and $g(x)$ in the MSSM is different. In fact, the former now reads [45]

$$f(x) \;=\; -x \left[\frac{2}{x^2-1} + \ln\left(1 + \frac{1}{x^2}\right) \right] . \tag{134}$$

Comparing its behavior to the loop function in (59), relevant in the SM, we notice the following: both of them can be zero, however, the exact position is slightly different ($x \approx 0.42$ for the SM loop function and $x \approx 0.34$ for $f(x)$ in (134)); otherwise, their values are of the same order of magnitude for $x \lesssim 1$; for $x \gtrsim 1$ it holds to good approximation that $(f(x))_{\rm SUSY} \approx 2\,(f(x))_{\rm SM}$. They have a divergence at $x = 1$ in common. Concerning the loop function $g(x)$ which enters in the flavored CP asymmetries, we have the exact relation $(g(x))_{\rm SUSY} = 2\,(g(x))_{\rm SM}$, where $(g(x))_{\rm SM}$ is given in (122).

At the same time, additional washout effects are induced by the SUSY particles. Also the numerical factor in (57) slightly changes and reads 1.48×10^{-3} due to the additional particles. Also the sphaleron conversion factor changes due to the presence of a second Higgs doublet.

Summarizing all the effects and assuming thermal leptogenesis to be the dominant mechanism for generating a lepton (and thus the baryon) asymmetry, the results for the baryon asymmetry of the Universe $Y_{B,\rm SUSY}$ in the SUSY framework are obtained by rescaling $Y_{B,\rm SM}$, computed in a SM context, in the following way [49]

$$\begin{aligned} & Y_{B,\rm SUSY} \approx \sqrt{2}\, Y_{B,\rm SM} \quad \text{in the regime of strong washout} \\ \text{and} \qquad & Y_{B,\rm SUSY} \approx 2\sqrt{2}\, Y_{B,\rm SM} \quad \text{in the regime of weak washout.} \end{aligned} \tag{135}$$

6 $0\nu\beta\beta$ decay

In this section we study the impact of leptonic CP violation, focussing on $0\nu\beta\beta$ decay of even-even nuclei. This process, which has not been observed so far, is important to test the Majorana nature of neutrinos and it explicitly depends on the values of the Majorana phases α and β (see e.g. [62]

[15] For such values of $\tan\beta$ also the suppression of the bottom quark mass with respect to the top quark mass has to originate from the flavor symmetry. Thus, we expect in concrete models that also the down type quark masses stem from non-renormalizable operators only.

for a recent review). Therefore, it is phenomenologically relevant in order to put constraints on the scenarios introduced in subsection 2.2. Earlier studies of $0\nu\beta\beta$ decay in the context of models with combined flavor and CP symmetries can be found in [26, 39] as well as the last reference in [23] and the first reference in [28]. After a short subsection on some general information about the quantity measurable in $0\nu\beta\beta$ decay we discuss predictions for it in some of the scenarios of mixing separately, referring in particular to the examples for which leptogenesis has been analyzed.

6.1 Preliminaries and general results

The half-life $T_{1/2}^{0\nu\beta\beta}$ of a decaying nuclear isotope via this process is

$$[T_{1/2}^{0\nu\beta\beta}]^{-1} \;=\; G^{0\nu} \left|M^{0\nu}\right|^2 \frac{m_{ee}^2}{m_e^2}\,, \tag{136}$$

where $G^{0\nu}$ is a phase space factor, $M^{0\nu}$ is the nuclear matrix element (NME) for a lepton number violating transition, m_e is the electron mass and m_{ee} is the effective Majorana neutrino mass. The values of $G^{0\nu}$ and $M^{0\nu}$ can be computed and depend on the nuclear isotope, whereas m_{ee} is expressed only in terms of neutrino masses and lepton mixing parameters,

$$m_{ee} = \left|U_{PMNS,11}^2\, m_1 + U_{PMNS,12}^2\, m_2 + U_{PMNS,13}^2\, m_3\right|\,, \tag{137}$$

that, according to the parametrization of U_{PMNS} reported in appendix A.1, reads

$$m_{ee} = \left|\cos^2\theta_{12}\,\cos^2\theta_{13}\, m_1 + \sin^2\theta_{12}\,\cos^2\theta_{13}\, e^{i\alpha}\, m_2 + \sin^2\theta_{13}\, e^{i\beta}\, m_3\right|\,. \tag{138}$$

An upper bound on the effective Majorana neutrino mass has been recently determined by several experiments, each one designed for a specific nuclear isotope: GERDA (^{76}Ge) [63], KamLAND-Zen (^{136}Xe) [64], EXO-200 (^{136}Xe) [65], CUORE-0 (^{130}Te) [66], and NEMO 3 (^{100}Mo) [67]. The strongest bound on m_{ee} is set by the KamLAND-Zen experiment

$$m_{ee} \;<\; (0.14 \div 0.28)\ \text{eV} \quad \text{at } 90\%\ \text{C.L.} \tag{139}$$

with the largest uncertainty coming from the computation of the associated NME.

For a hierarchical light neutrino mass spectrum, i.e. for $m_0 \approx 0$ in (19) and (20), the expected value of m_{ee} strongly depends on the ordering of the neutrino masses. In fact, in this limit we have from (138), for NO

$$m_{ee}^{\rm NO} \approx \left|\sin^2\theta_{12}\,\cos^2\theta_{13}\, e^{i\alpha}\,\sqrt{\Delta m^2_{\rm sol}} + \sin^2\theta_{13}\, e^{i\beta}\,\sqrt{\Delta m^2_{\rm atm}}\right| \tag{140}$$

and for IO

$$m_{ee}^{\rm IO} \approx \left|\cos^2\theta_{12} + \sin^2\theta_{12}\, e^{i\alpha}\right|\,\cos^2\theta_{13}\,\sqrt{|\Delta m^2_{\rm atm}|}\,. \tag{141}$$

In the last expression we have neglected the subdominant term proportional to $\Delta m^2_{\rm sol}$. For a given value of θ_{12} and θ_{13}, the maximum and minimum of m_{ee} are obtained for trivial Majorana phases, see e.g. the lower-left panel of figure 9. In particular, for IO, using that $\sin^2\theta_{12} \approx 1/3$, we have

$$\frac{1}{3}\,\sqrt{|\Delta m^2_{\rm atm}|} \lesssim m_{ee}^{\rm IO} \lesssim \sqrt{|\Delta m^2_{\rm atm}|}\,. \tag{142}$$

In the case of a QD spectrum, we expect from (138) m_{ee} to be proportional to m_0, for both NO and IO. Indeed, neglecting $\sin\theta_{13}$, we have

$$m_{ee}^{\rm QD} \approx \sqrt{1 - \sin^2\frac{\alpha}{2}\,\sin^2 2\theta_{12}}\;\; m_0\,, \tag{143}$$

which, for $\sin^2\theta_{12} \approx 1/3$ and a trivial Majorana phase α, takes values in the interval

$$\frac{1}{3}\, m_0 \lesssim m_{ee}^{\rm QD} \lesssim m_0\,. \tag{144}$$

As is well-known, for NO m_{ee} can be strongly suppressed due to cancellations for m_0 between 10^{-3} eV and 0.01 eV. This can occur in principle, if both Majorana phases are trivial or both are non-trivial. In our numerical analysis we find examples for both situations, see e.g. case 1) for the former one (compare (152)) and case 3 a) for the latter one, see figure 10.

In figures 8-10 we also show in light-blue (orange) the most general region in the $m_0 - m_{ee}$ plane for NO (IO), obtained by varying the lepton mixing angles within the experimentally preferred $3\,\sigma$ ranges, see appendix A.2, and all CP varies between 0 and $2\,\pi$. The borders of the areas are associated to trivial Majorana phases α and β and do depend on the lower and upper $3\,\sigma$ limit of the the solar mixing angle θ_{12}. These theoretically allowed areas are constrained by the current strongest upper limit in (139) on the effective Majorana neutrino mass given by the KamLAND-Zen collaboration as well as the bound on the lightest neutrino mass given by the Planck collaboration, see (23). The former is displayed as horizontal dashed line in figures 8-10, while the latter as vertical one.

6.2 Predictions of m_{ee} in case 1)

In this scenario the Majorana phase β (as well as the Dirac phase δ) is always trivial, for any choice of θ and the group theoretical parameters n and s (or their combination ϕ_s), while α can take non-trivial values, see (35). Then, for a hierarchical neutrino mass spectrum ($m_0 \approx 0$), m_{ee} reads

$$m_{ee}^{\rm NO} \;\approx\; \frac{1}{3}\left|\sqrt{\Delta m^2_{\rm sol}} + 2\,(-1)^{k_1+k_2}\,\sin^2\theta\, e^{6\,i\,\phi_s}\,\sqrt{\Delta m^2_{\rm atm}}\right|\,, \tag{145}$$

$$m_{ee}^{\rm IO} \;\approx\; \frac{1}{3}\left|1 + 2\,(-1)^{k_1}\, e^{6\,i\,\phi_s}\,\cos^2\theta\right|\,\sqrt{|\Delta m^2_{\rm atm}|}\,. \tag{146}$$

Remembering that θ is close to 0 or π, $m_{ee}^{\rm IO}$ can be simplified to

$$m_{ee}^{\rm IO} \;\approx\; \frac{1}{3}\,\sqrt{5 \pm 4\,\cos 6\,\phi_s}\,\sqrt{|\Delta m^2_{\rm atm}|}\,, \tag{147}$$

with the plus (minus) sign corresponding to even (odd) k_1. Similarly, we obtain for a QD spectrum

$$m_{ee}^{\rm QD} \;\approx\; \frac{1}{3}\left|(-1)^{k_1} + 2\,e^{6\,i\,\phi_s}\,\left(\cos^2\theta + (-1)^{k_2}\,\sin^2\theta\right)\right|\,m_0\,. \tag{148}$$

As one can see, for an even k_2 the effective Majorana neutrino mass is independent of θ

$$m_{ee}^{\rm QD} \;\approx\; \frac{1}{3}\,\sqrt{5 \pm 4\,\cos 6\,\phi_s}\;\,m_0\,, \tag{149}$$

with the plus (minus) sign valid for even (odd) k_1. Since $\theta \approx 0,\, \pi$, (149) is a good approximation also in the case of k_2 being odd. This result is consistent with the one found in (143) for the generic case.

In the case $n = 4$, that is the smallest value of the group theoretical parameter n allowing for non-trivial α (see table 1), we obtain for $s = 0, 2$ ($s = 1, 3$) and k_2 even

$$m_{ee}^{\rm NO} \approx 0.0350\;(0.0323)\;\text{eV} \quad \text{and} \quad m_{ee}^{\rm IO} \approx 0.0175\;(0.0165)\;\text{eV} \tag{150}$$

using the best fit values of the neutrino squared mass differences, see (21), $m_0 = 10^{-4}$ eV, and $\theta \approx 0.18$ or $\theta \approx 2.96$ that lead to the best fitting of the experimental data on mixing angles, see [27]. Similarly, for a QD spectrum, we find for $m_0 = 0.1$ eV, $s = 0, 2$ ($s = 1, 3$) and k_2 even

$$m_{ee}^{\rm QD} \approx 0.108\;(0.0814)\;\text{eV}\,. \tag{151}$$



Figure 8: **Case 2).** Effective Majorana neutrino mass as a function of the lightest neutrino mass, for $n = 10$, $u = 4$ and two values of v. The parameter θ is varied in the range $1.40 \lesssim \theta \lesssim 1.44$. The neutrino mass squared differences are randomly chosen within their 3σ intervals. The blue (red) region corresponds to the variation of m_{ee} in the case of NO (IO), for all the possible combinations of $k_{1,2}$. The particular choice $k_1 = k_2 = 0$ is highlighted by the dark-grey areas. The plot on the right panel also gives the expected m_{ee} for $v = 24$ and $1.70 \lesssim \theta \lesssim 1.74$, with the dark-grey area equal to $k_1 = 1$ and $k_2 = 0$. The other allowed values of v ($v = 12$ and $v = 18$) give predictions similar to the ones shown here. The light-blue and orange areas are the expectations obtained from the general expression of m_{ee} (137) for NO and IO, respectively, where the neutrino oscillation parameters are taken within the 3σ intervals reported in appendix A.2. The borders of these areas are obtained for CP conserving Majorana phases α and β and depend on the 3σ range of the solar mixing angle.

Smaller values of m_{ee} result for the same m_0, but k_2 odd, if the neutrino spectrum is either hierarchical with NO or QD due to cancellations in the expressions in (145) and (148). Since $m_{ee}^{\rm IO}$ does not depend on k_2, see (147), there is no such effect for the case of a hierarchical inversely ordered neutrino mass spectrum.

As in the generic case, for trivial α occurring for $s = 0,\, 2$, m_{ee} is strongly suppressed for certain values of the lightest neutrino mass in case of NO. In particular, $m_{ee}^{\rm NO}$ is smaller than 10^{-4} eV for

$$0.0023 \text{ eV} \lesssim m_0 \lesssim 0.0028 \text{ eV} \quad \text{and} \quad 0.0066 \text{ eV} \lesssim m_0 \lesssim 0.0087 \text{ eV} \tag{152}$$

with the range of the intervals resulting from the variation of the neutrino mass squared differences in their $3\,\sigma$ range and $0.168 \lesssim \theta \lesssim 0.195$ or $2.95 \lesssim \theta \lesssim 2.97$. A cancellation for m_0 being the first interval requires k_1 odd and k_2 even, while m_0 in the second interval k_1 and k_2 odd. We note that due to the constraints on mixing angles and CP phases the range of m_0 in which m_{ee} can be very small for NO is considerably reduced, especially, the fact that the smallest of m_0 for which this cancellation can occur is larger than in the general case results from the lower bound on the solar mixing angle $\sin^2\theta_{12} \gtrsim 1/3$.

6.3 Predictions of m_{ee} in case 2)

In this case both Majorana phases α and β can have in general non-trivial values, depending on θ and the group theoretical parameters n, u and v (or their combinations ϕ_u and ϕ_v). Taking the lepton mixing matrix $U_{PMNS,2}$ as defined in (36), the general expression (137) for a hierarchical

spectrum can be approximated as

$$m_{ee}^{\rm NO} \approx \frac{1}{3}\left|\sqrt{\Delta m^2_{\rm sol}} - 2\,(-1)^{k_1+k_2}\,e^{i\,\phi_v}\,\left(\cos\theta\,\sin\frac{\phi_u}{2} - i\,\sin\theta\,\cos\frac{\phi_u}{2}\right)^2\,\sqrt{\Delta m^2_{\rm atm}}\right|\,, \quad (153)$$

$$m_{ee}^{\rm IO} \approx \frac{1}{3}\left|1 + (-1)^{k_1}\,e^{i\,\phi_v}\,\left(\cos\phi_u + \cos 2\theta - i\,\sin 2\theta\,\sin\phi_u\right)\right|\,\sqrt{|\Delta m^2_{\rm atm}|}\,. \quad (154)$$

For a QD spectrum and k_2 even, as in case 1), the resulting m_{ee} is actually independent of θ

$$m_{ee}^{\rm QD} \approx \frac{1}{3}\,\sqrt{3 + 2\,\cos 2\,\phi_u \pm 4\,\cos\phi_u\,\cos\phi_v}\;\,m_0\,, \quad (155)$$

with the upper (lower) sign standing for even (odd) k_1. For k_2 odd instead one can show that m_{ee} is independent of ϕ_u, if $\theta \approx 0,\ \pi$ or $\pi/2$ that are the typical values required to reproduce the observed lepton mixing angles, see [27],

$$m_{ee}^{\rm QD} \approx \frac{1}{3}\,\sqrt{5 \pm 4\,\cos\phi_v}\;\,m_0\,, \quad (156)$$

with again the plus (minus) sign referring to even (odd) values of k_1. This expression coincides with the one derived in the generic case in (143), using that in case 2) holds $\cos\alpha \approx (-1)^{k_1}\,\cos\phi_v$, similar to the approximate relation in (41).

Expressions similar to (153-156) are obtained, if we perform a permutation of the rows of $U_{PMNS,2}$ using (42). In this case m_{ee} can be computed by applying the transformations given in (44-45) to the approximations derived.

We note that the quantity m_{ee} is invariant under the set of transformations (see below (45))

$$\theta \to \pi - \theta\,, \quad v \to k\,n - v\ (k \text{ odd}) \quad \text{and} \quad k_1 \to k_1 + 1\,, \quad (157)$$

which shows that results for different values of v are related to each other, if we also take into account that the interval of θ has to be changed.

In the explicit example, we choose for the flavor group the same index $n = 10$ and as parameter characterizing the CP symmetry also $u = 4$, as in the numerical study of leptogenesis in subsection 3.4. We thus discuss a case in which we can use formulae shown above, only after having applied the transformations in (44). This scenario predicts $|\sin\beta| \approx 1$, while $|\sin\alpha|$ can take three different values, depending on the parameter v (or ϕ_v), i.e. $v = 0$, 6, 12, 18 and 24, see table 2. We display in figure 8 the prediction of m_{ee} as a function of the lightest neutrino mass m_0, for two different choices of v: $v = 0$ in the left and $v = 6$ in the right panel. These values lead either to $\alpha \approx 0,\ \pi$ ($v = 0$) or to almost maximal α ($v = 6$). The blue and red regions in each plot correspond to the predictions for NO and IO, respectively, in which we allow for all four possible combinations of k_1 and k_2. Among these, we mark in dark-grey the areas with $k_1 = k_2 = 0$. Moreover, we vary θ in the range $1.40 \lesssim \theta \lesssim 1.44$ and the neutrino mass squared differences within their 3σ intervals, see appendix A.2. Using (157) we see that for $v = 0$ values of θ in the second interval $1.70 \lesssim \theta \lesssim 1.74$ lead to the same allowed areas, up to exchange $k_1 = 0$ with $k_1 = 1$. For $v \neq 0$ instead, applying (157) shows, for example, that the plot on the right panel for $v = 6$ and $1.40 \lesssim \theta \lesssim 1.44$ is the same for $v = 24$ and θ in the second interval $1.70 \lesssim \theta \lesssim 1.74$. The only difference is that the dark-grey area corresponds for $v = 24$ to the choice $k_1 = 1$ and $k_2 = 0$. The predictions of m_{ee} for $v = 12$ and $v = 18$ are related to each other in a similar way. Figures for these two values of v resemble to the one displayed for $v = 0$.

The most important feature of this case is the fact that there is no cancellation in m_{ee} for any values of v in the case of NO. Furthermore, we note that for $v = 0$ the Majorana phases are nearly trivial which explains why the predictions for IO coincide with the borders of the area allowed in the generic case. Using (156) with $v = 0$ shows that $m_{ee}^{\rm QD}$ either equals $m_{ee}^{\rm QD} \approx m_0$ for k_1 odd or $m_{ee}^{\rm QD} \approx m_0/3$ for k_1 even (remember k_1 has to be replaced by $k_1 + 1$ in (156) in order to apply to the case at hand). For m_0 small and NO instead m_{ee} takes values in the interval



Figure 9: **Case 3 b.1).** m_{ee} versus m_0 for $n = 8$, $m = 4$ and three possible choices of s. For a given s, θ takes values in the interval reported in table 3. In all the plots the mass squared differences are varied within 3σ (see appendix A.2). The blue and red areas correspond to the NO and IO spectrum, respectively, for any combination of $k_{1,2}$. In dark-grey it is displayed the region with $k_1 = k_2 = 0$. The light-blue and orange areas are the same as shown in figure 8. Results for $s > n/2$ are obtained from the ones presented by making use of the transformation in (54).

0.0026 eV $\lesssim m_{ee}^{\rm NO} \lesssim$ 0.0035 eV. In constrast, for $v = 6$ we see two different regimes realized: $m_{ee}^{\rm NO} \approx 0.0018$ eV for $k_1 + k_2$ odd and $m_{ee}^{\rm NO} \approx 0.0039$ eV for $k_1 + k_2$ even, respectively. Since the Majorana phase α is non-trivial and not small for $v = 6$, we find a non-trivial lower bound on m_{ee} in the case of IO that is by a factor of two larger than the generic lower bound, $m_{ee}^{\rm IO} \approx 0.031$ eV. The other value of m_{ee} is $m_{ee}^{\rm IO} \approx 0.039$ eV, arising, if a k_1 is even. Two different values are also obtained in the regime of QD neutrino masses where we predict for $m_0 = 0.1$ eV, according to (156), $m_{ee}^{\rm QD} \approx 0.065$ eV or $m_{ee}^{\rm QD} \approx 0.083$ eV depending on the value of k_1. Future experiments searching for $0\nu\beta\beta$ decay [68–70] will probe almost all the IO orange region, up to $m_{ee} \approx 0.02$ eV, thus allowing to distinguish between the different CP violating scenarios.

6.4 Predictions of m_{ee} in case 3)

We would like to discuss the effective Majorana neutrino mass given in the last case introduced in subsection 2.2. We first consider case 3 b.1) that is characterized by the lepton mixing matrix in (46). We focus on the choice $m = n/2$, as done in our analysis of leptogenesis. In this case, it follows directly from (55) that m_{ee} must be invariant under the replacement of θ with $\pi - \theta$.

Again, we can derive simple approximations that work well for the hierarchical and QD regimes.

Thus, for vanishing m_0 we have

$$m_{ee}^{\rm NO} \approx \frac{1}{3}\left|\sin^2\theta\sqrt{\Delta m^2_{\rm sol}} + (-1)^{k_1}\,\cos^2\theta\sqrt{\Delta m^2_{\rm atm}}\right|\,, \tag{158}$$

$$m_{ee}^{\rm IO} \approx \frac{1}{3}\left|2 + (-1)^{k_2}\,e^{6\,i\,\phi_s}\sin^2\theta\right|\,\sqrt{|\Delta m^2_{\rm atm}|}\,. \tag{159}$$

Interestingly enough, $m_{ee}^{\rm NO}$ is independent of ϕ_s (and thus the specific CP transformation) and it takes two distinct values for k_1 even and odd, respectively. Using for the neutrino mass squared differences the best fit values and choosing $\theta \approx 1.31$ (which sets the reactor angle at the experimental best fit value [27]), we have

$$m_{ee}^{\rm NO} \;=\; 0.0038\;(0.0016)\;\;{\rm eV}\,, \qquad \text{for an even (odd)}\;\; k_1\;\; \text{and} \tag{160}$$

$$m_{ee}^{\rm IO} \;=\; \left|0.015\,e^{6\,i\,\phi_s} \,+\, (-1)^{k_2}\,0.033\right|\;\;{\rm eV}\,. \tag{161}$$

In contrast, for the QD spectrum we obtain

$$m_{ee}^{\rm QD} \approx \frac{1}{3}\left|2 \,+\, (-1)^{k_2}\,e^{6\,i\,\phi_s}\left(\sin^2\theta \,+\, (-1)^{k_1}\cos^2\theta\right)\right|\,m_0\,, \tag{162}$$

which becomes independent of θ for an even k_1. Using $\theta \approx \pi/2$ we can further simplify (162) and obtain the expression in (149), which is valid in case 1).

We show in figure 9, the expected m_{ee} versus m_0 for the values of θ and the group theoretical parameters reported in table 3. The blue (red) area corresponds to NO (IO), for any combination of $k_{1,2}$, with the particular case $k_1 = k_2 = 0$ depicted in dark-grey. The light-blue and the orange regions in each plot are the same as those shown in figure 8. The experimental limits on m_{ee} and m_0 in the plots (horizontal and vertical dashed lines) are given in (139) and (23).

For $s = 1$ and $s = 2$, the Majorana phases are indeed non-trivial (see table 3) and m_{ee} has a lower bound in the NO spectrum. This is similar to what we found in the numerical example of case 2), discussed above. In the case of IO we have that for $s = 2$ ($\phi_s = \pi/4$), $m_{ee}^{\rm IO}$ is actually independent of $k_{1,2}$, see (159), and we have only one (in this case dark-grey) narrow shaded area within the orange region of the right panel, which corresponds to $m_{ee}^{\rm IO} \approx 0.036$ eV. Instead, for $s = 1$ we have in the hierarchical regime $m_{ee}^{\rm IO} \approx 0.024$ (0.045) eV for even (odd) k_2, compare (161). Similarly, we find for QD neutrino mass spectrum two possible values for $s = 1$, $m_{ee} \approx 0.49\,m_0$ and $m_{ee} \approx 0.93\,m_0$, whereas for $s = 2$ we only find $m_{ee} \approx 0.75\,m_0$. In addition, one can show that, in the case of NO and $s = 2$, m_{ee} only depends on k_1 in all the range of m_0. This explains why for this choice of s we have only two distinct regions in the hierarchical regime. We have for $m_0 = 10^{-4}$ eV and $s = 1$ that $m_{ee}^{\rm NO}$ varies in the interval 0.0034 eV $\lesssim m_{ee}^{\rm NO} \lesssim$ 0.0040 eV (0.0012 eV $\lesssim m_{ee}^{\rm NO} \lesssim$ 0.0020 eV) for even (odd) k_1, see (158). For the same m_0 and $s = 2$ we find, as expected, very similar intervals, that is 0.0035 eV $\lesssim m_{ee}^{\rm NO} \lesssim$ 0.0039 eV (0.0015 eV $\lesssim m_{ee}^{\rm NO} \lesssim$ 0.0019 eV) for even (odd) k_1. In both cases these results are in agreement with the analytic estimates for $m_{ee}^{\rm NO}$ given in (160).

In contrast, in the case $s = 4$ the Majorana phases α and β are trivial and the effective Majorana neutrino mass can be strongly suppressed in the NO hierarchical regime, as shown in bottom-left panel of figure 9. Notice that our approach constrains not only the CP phases but also the mixing angles to take values in narrow ranges, see table 3. Therefore, the suppression of m_{ee} occurs only for two small intervals of m_0, which depend on the integers $k_{1,2}$: $m_{ee}^{\rm NO} \lesssim 10^{-4}$ for 0.0019 eV $\lesssim m_0 \lesssim$ 0.0033 eV (0.0059 eV $\lesssim m_0 \lesssim$ 0.0074 eV) and $k_1 = k_2 = 1$ ($k_1 = k_2 = 0$). In the strongly hierarchical regime, that is for $m_0 \lesssim 10^{-4}$ eV, we have, instead, 0.0035 eV $\lesssim m_{ee}^{\rm NO} \lesssim$ 0.0039 eV (0.0012 eV $\lesssim m_{ee}^{\rm NO} \lesssim$ 0.0020 eV) for even (odd) k_1, in agreement with the estimate in (160).

Finally, we remark that for case 3a), which requires the group index n to be larger than 10, several choices of the CP transformation X (or, equivalently, the parameter s) are admitted. Some of these allow to have strong cancellations in the expression of m_{ee}, while both Majorana phases are non-trivial. One example is considered in figure 10, where it is displayed the prediction of m_{ee} for $n = 16$, $m = 1$ and $s = 2$ (left panel) and $s = 3$ (right panel). The parameter θ is taken in the



Figure 10: **Case 3a).** The same as in figure 9 for $n = 16$, $m = 1$ and two choices of s. The parameter θ is varied in the intervals given in the text. For these values of s at least one of the Majorana phases is non-trivial and strong cancellations in m_{ee} are realized in the case of NO.

interval $2.28 \lesssim \theta \lesssim 2.56$ ($0 \lesssim \theta \lesssim 0.540$) for $s = 2$ ($s = 3$). For $k_1 = k_2 = 0$ the Majorana phases α and β are constrained to lie in the intervals: $0.330 \lesssim \sin\alpha \lesssim 0.712$ ($-0.392 \lesssim \sin\alpha \lesssim 0.383$) and $0.654 \lesssim \sin\beta \lesssim 0.872$ ($-0.675 \lesssim \sin\beta \lesssim 0$) for $s = 2$ ($s = 3$). Consequently, strong cancellations in m_{ee} for NO occur, if $s = 2$ and

$$0.0043 \text{ eV} \lesssim m_0 \lesssim 0.0052 \text{ eV} \quad \text{for } k_1 = k_2 = 1 \tag{163}$$

or $s = 3$ and

$$0.0019 \text{ eV} \lesssim m_0 \lesssim 0.0027 \text{ eV} \quad \text{for } k_1 = k_2 = 0 \tag{164}$$

$$\text{and} \qquad 0.0062 \text{ eV} \lesssim m_0 \lesssim 0.0072 \text{ eV} \quad \text{for } k_1 = k_2 = 1\,, \tag{165}$$

as one can see in figure 10. Such strong cancellations can also be achieved for $s = 0$ and $s = 8$ where a common accidental CP symmetry in the charged lepton and neutrino sector is present [27]. For the other values of $s \leq 8$, not shown here, the resulting m_{ee} has a lower bound, typically $m_{ee} \gtrsim 10^{-3}$ eV, within the range of m_0 shown in the figures.

In summary, we have shown that in the case of NO all three different situations can be realized: no cancellations in m_{ee} for case 2), $n = 10$ and $u = 4$, strong suppression of m_{ee} and trivial Majorana phases for case 3b.1), $n = 8$, $m = 4$ and $s = 4$, as well as cancellations in the presence of two non-trivial Majorana phases for case 3a), $n = 16$, $m = 1$ and $s = 2$ or $s = 3$. For IO and QD we observe that m_{ee} can only obtain values in a very limited range for all possible m_0, which mainly depends on the chosen CP transformation and the parameters k_1 and k_2. For this reason, at least part of these scenarios, see (149), (156) and (162), will be testable in near future experiments looking for $0\nu\beta\beta$ decay.

7 Summary

We have studied leptogenesis and $0\nu\beta\beta$ decay in a scenario with a flavor G_f and a CP symmetry that are broken non-trivially in the charged lepton and neutrino sectors. We have chosen as G_f groups belonging to the series $\Delta(3\,n^2)$ and $\Delta(6\,n^2)$, $n \geq 2$, while the CP symmetry is represented

by a CP transformation X that is a unitary and symmetric matrix, acting non-trivially in flavor space. The residual symmetry G_e in the charged lepton sector is fixed to a Z_3 subgroup of G_f and G_ν, the symmetry preserved in the neutrino sector, is given by the direct product $Z_2 \times CP$ with $Z_2 \subset G_f$. In this way, the charged lepton mass matrix can always be constrained to be diagonal, while the light neutrino mass matrix contains four independent parameters, corresponding to the three neutrino masses and $\theta \in [0, \pi)$, on which lepton mixing angles and CP phases depend.

Under the assumption of three RH neutrinos and a Dirac Yukawa coupling Y_D that is invariant under G_f and CP the CP asymmetries $\epsilon_i^{(\alpha)}$ vanish in the case of flavored as well as unflavored leptogenesis. If Y_D is taken to be invariant under the residual symmetry $G_\nu = Z_2 \times CP$ only, ϵ_i^α become non-vanishing and a non-zero value of Y_B can be achieved via flavored leptogenesis. Still, in the case of unflavored leptogenesis this is not sufficient and ϵ_i turn out to vanish.

In our study of unflavored leptogenesis we introduce corrections δY_D to Y_D in order to obtain ϵ_i non-zero. These are expected to be in general proportional to the symmetry breaking parameter κ of our scenario. A particularly interesting case is to assume that δY_D is induced by the breaking of G_f to G_e and thus is invariant under G_e, the residual symmetry of the charged lepton sector. The two main consequences are: the suppression of the CP asymmetries $\epsilon_i \propto \kappa^2$ and the fact that the Dirac and Majorana phases, potentially measurable in terrestrial experiments, determine the sign of Y_B. The first observation has already been made in approaches with a flavor symmetry only, whereas the second one, namely the prediction of the sign of Y_B, is only possible thanks to the presence of a CP symmetry that controls the CP phases. Especially, we have found the following: phases that are present in the correction δY_D are irrelevant for the sign of Y_B at LO; if not suppressed (due to a special choice of the neutrino masses), we expect the terms involving the sine of the Majorana phase α to dominate the CP asymmetries and thus the sign of Y_B (see figures 4 and 5, the upper-right and lower panel in figure 6 and the upper two panels in figure 7); in case the dominant contribution to the CP asymmetries arises from terms involving the Dirac phase the sign of Y_B cannot be predicted and positive and negative values are equally possible (see the upper-left panel of figure 6 and the lower panel of figure 7). These features are in addition confirmed by our analysis of a scenario in which the lepton mixing is of a general form (and not constrained by symmetries).

For flavored leptogenesis we have demonstrated with several examples that the sign of the CP asymmetries ϵ_i^α depends on additional input, e.g. the relative size of the parameters of the correction δY_D, and thus fixing the sign of ϵ_i^α is in general impossible, having only information about the CP phases and the light neutrino mass ordering.

We have argued that the symmetry breaking scenario considered by us is well-motivated and have presented examples of non- as well as SUSY realizations of this scenario in appendix D where we show that small corrections δY_D of the advocated form are naturally obtained.

We have also studied in detail the predictions for m_{ee}, accessible in experiments searching for $0\nu\beta\beta$ decay. The following interesting properties are found: the constraints on CP phases and lepton mixing angles allow for cases in which no cancellation in m_{ee} occurs even for normally ordered light neutrino masses and thus $m_{ee} \gtrsim 0.002\,\text{eV}$ can be achieved; in the case of an inverted neutrino mass ordering such constraints can lead to values of $m_{ee} \gtrsim 0.05\,\text{eV}$ thus increasing the chances to measure this quantity in the not-too-far future.

In summary, we have thoroughly analyzed a scenario in which flavor and CP symmetries can determine low and high energy CP phases together with the lepton mixing angles. Thus, phenomena requiring CP violation can be related. In particular, we have shown that the knowledge of low energy CP phases (and the light neutrino mass spectrum) can be sufficient for fixing the sign of the baryon asymmetry of the Universe, if the latter is generated via unflavored thermal leptogenesis.

Acknowledgements

C.H. and E.M. thank the he Excellence Cluster 'Universe' at the Technische Universität München for partial support during the preparation of this work. We notice that important parts of the final results of this work were presented by E.M. at the workshop "Nu@Fermilab", 21-25 July 2015, in

Fermilab, Chicago, and by C.H. at the weekly theory seminar of the high energy physics group in Southampton on 29 January 2016.

A Conventions for mixing angles and CP invariants, and global fit results

In this appendix we fix our conventions of mixing angles and of the CP invariants J_{CP}, I_1 and I_2 and list the latest global fit results [14].

A.1 Conventions for mixing angles and CP invariants

As parametrization of the PMNS mixing matrix we use

$$U_{PMNS} = \tilde{U} \text{diag}(1, e^{i\alpha/2}, e^{i(\beta/2+\delta)}) \,, \tag{166}$$

with $\tilde{U}$ being of the form of the Cabibbo-Kobayashi-Maskawa (CKM) matrix V_{CKM} [71]

$$\tilde{U} = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta} & c_{23}c_{13} \end{pmatrix} \tag{167}$$

and $s_{ij} = \sin\theta_{ij}$ and $c_{ij} = \cos\theta_{ij}$. The mixing angles θ_{ij} range from 0 to $\pi/2$, while the Majorana phases α, β as well as the Dirac phase δ take values between 0 and 2π. The Jarlskog invariant J_{CP} reads [72]

$$\begin{aligned} J_{CP} &= \text{Im}\left[U_{PMNS,11} U^*_{PMNS,13} U^*_{PMNS,31} U_{PMNS,33}\right] \\ &= \frac{1}{8} \sin 2\theta_{12} \sin 2\theta_{23} \sin 2\theta_{13} \cos\theta_{13} \sin\delta \,. \end{aligned} \tag{168}$$

Similar invariants, called I_1 and I_2, can be defined which depend on the Majorana phases α and β, see e.g. [73]

$$I_1 = \text{Im}[U^2_{PMNS,12}(U^*_{PMNS,11})^2] = s^2_{12}c^2_{12}c^4_{13}\sin\alpha \,, \tag{169}$$

$$I_2 = \text{Im}[U^2_{PMNS,13}(U^*_{PMNS,11})^2] = s^2_{13}c^2_{12}c^2_{13}\sin\beta \,. \tag{170}$$

A third CP invariant related to the Majorana phases can be introduced

$$I_3 = \text{Im}[U^2_{PMNS,13}(U^*_{PMNS,12})^2] = c^2_{13}s^2_{13}s^2_{12}\sin(\beta - \alpha) \,. \tag{171}$$

Notice that the Dirac phase has a physical meaning only if all mixing angles are different from 0 and $\pi/2$, as indicated by the data. Analogously, the vanishing of the invariants $I_{1,2}$ only implies $\sin\alpha = 0$, $\sin\beta = 0$, if solutions with $\sin 2\theta_{12} = 0$, $\cos\theta_{13} = 0$ or $\sin 2\theta_{13} = 0$, $\cos\theta_{12} = 0$ are discarded. Furthermore, notice that one of the Majorana phases becomes unphysical, if the lightest neutrino mass vanishes.

A.2 Global fit results

We use in our numerical analysis the results of mixing angles, the CP phase δ and the mass squared differences Δm^2_{sol} and Δm^2_{atm} taken from [14]. For NO the best fit values of $\sin^2\theta_{ij}$ and δ (given here in radian), the $1\,\sigma$ errors as well as $3\,\sigma$ ranges are

$$\begin{aligned} &\sin^2\theta_{13} = 0.0218^{+0.0010}_{-0.0010} && \text{and} && 0.0186 \leq \sin^2\theta_{13} \leq 0.0250 \\ &\sin^2\theta_{12} = 0.304^{+0.013}_{-0.012} && \text{and} && 0.270 \leq \sin^2\theta_{12} \leq 0.344 \\ &\sin^2\theta_{23} = 0.452^{+0.052}_{-0.028} && \text{and} && 0.382 \leq \sin^2\theta_{23} \leq 0.643 \\ &\delta = 5.34^{+0.68}_{-1.22} && \text{and} && 0 \leq \delta \leq 2\,\pi \end{aligned} \tag{172}$$

as well as for the mass squared differences $\Delta m^2_{\rm sol}$ and $\Delta m^2_{\rm atm}$

$$\Delta m^2_{\rm sol} = \left(7.50^{+0.19}_{-0.17}\right) \times 10^{-5}\,{\rm eV}^2 \quad \text{and} \quad \Delta m^2_{\rm atm} = \left(2.457^{+0.047}_{-0.047}\right) \times 10^{-3}\,{\rm eV}^2 \tag{173}$$

and

$$\begin{aligned} 7.03 \times 10^{-5}\,{\rm eV}^2 \leq \quad & \Delta m^2_{\rm sol} \quad \leq 8.03 \times 10^{-5}\,{\rm eV}^2 \\ 2.317 \times 10^{-3}\,{\rm eV}^2 \leq \quad & \Delta m^2_{\rm atm} \quad \leq 2.607 \times 10^{-3}\,{\rm eV}^2 \,. \end{aligned} \tag{174}$$

For IO instead the global fit analysis in [14] yields

$$\begin{aligned} &\sin^2\theta_{13} = 0.0219^{+0.0011}_{-0.0010} && \text{and} && 0.0188 \leq \sin^2\theta_{13} \leq 0.0251 \\ &\sin^2\theta_{12} = 0.304^{+0.013}_{-0.012} && \text{and} && 0.270 \leq \sin^2\theta_{12} \leq 0.344 \\ &\sin^2\theta_{23} = 0.579^{+0.025}_{-0.037} && \text{and} && 0.389 \leq \sin^2\theta_{23} \leq 0.644 \\ &\delta = 4.43^{+1.10}_{-1.08} && \text{and} && 0 \leq \delta \leq 2\,\pi \end{aligned} \tag{175}$$

as well as the mass squared differences

$$\Delta m^2_{\rm sol} = \left(7.50^{+0.19}_{-0.17}\right) \times 10^{-5}\,{\rm eV}^2 \quad \text{and} \quad \Delta m^2_{\rm atm} = \left(-2.449^{+0.048}_{-0.047}\right) \times 10^{-3}\,{\rm eV}^2 \tag{176}$$

and

$$\begin{aligned} 7.02 \times 10^{-5}\,{\rm eV}^2 \leq \quad & \Delta m^2_{\rm sol} \quad \leq 8.09 \times 10^{-5}\,{\rm eV}^2 \\ -2.590 \times 10^{-3}\,{\rm eV}^2 \leq \quad & \Delta m^2_{\rm atm} \quad \leq -2.307 \times 10^{-3}\,{\rm eV}^2 \,. \end{aligned} \tag{177}$$

B Generators of $\Delta(3\,n^2)$ and $\Delta(6\,n^2)$

Here we present the generators of the groups $\Delta(3\,n^2)$ and $\Delta(6\,n^2)$ that have been used in [27]. For further details on the groups $\Delta(3\,n^2)$ and $\Delta(6\,n^2)$ see [74] and [75], respectively. The generators a and c of $\Delta(3\,n^2)$ for the representation **3** that is faithful and irreducible for all groups can be chosen as

$$\begin{aligned} a &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & \omega & 0 \\ 0 & 0 & \omega^2 \end{pmatrix} \,, \\ c &= \frac{1}{3} \begin{pmatrix} 1+2\cos\phi_n & 1-\cos\phi_n-\sqrt{3}\sin\phi_n & 1-\cos\phi_n+\sqrt{3}\sin\phi_n \\ 1-\cos\phi_n+\sqrt{3}\sin\phi_n & 1+2\cos\phi_n & 1-\cos\phi_n-\sqrt{3}\sin\phi_n \\ 1-\cos\phi_n-\sqrt{3}\sin\phi_n & 1-\cos\phi_n+\sqrt{3}\sin\phi_n & 1+2\cos\phi_n \end{pmatrix} \end{aligned} \tag{178}$$

with $\omega = e^{\frac{2\pi i}{3}}$ and $\phi_n = \frac{2\pi}{n}$. They fulfill together with the generator $d = a^2 c\, a$ the relations

$$\begin{aligned} & a^3 = e \;\;,\;\; c^n = e \;\;,\;\; d^n = e \;\;, \\ & c\,d = d\,c \;\;,\;\; a\,c\,a^{-1} = c^{-1} d^{-1} \;\;,\;\; a\,d\,a^{-1} = c \end{aligned} \tag{179}$$

with e being the neutral element of the group. In order to arrive at the group $\Delta(6\,n^2)$ we have to add another generator b, chosen for the representation **3** as

$$b = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & \omega^2 \\ 0 & \omega & 0 \end{pmatrix} \tag{180}$$

that fulfills the following relations

$$b^2 = e \;\;,\;\; (a\,b)^2 = e \;\;,\;\; b\,c\,b^{-1} = d^{-1} \;\;,\;\; b\,d\,b^{-1} = c^{-1} \,. \tag{181}$$

We note that all groups have a trivial singlet **1** for which all elements of the group are represented by the unit.

C Results for CP asymmetries ϵ_i in the limit $z_1 = 0$

For completeness, we report here the formulae for the CP asymmetries ϵ_i that are obtained in the limit $z_1 = 0$ (equivalent to $\zeta = \pi/2,\, 3\pi/2$) and for only one non-vanishing CP phase.

First, we consider the case in which the Majorana phase α and the phase combination $\beta + 2\,\delta$ are trivial, i.e.

$$\alpha = k_\alpha\,\pi \;\;,\;\; \beta + 2\delta = k_\beta\,\pi \quad \text{with} \quad k_{\alpha,\beta} = 0, 1\;. \tag{182}$$

This choice takes our non-standard definition of the second Majorana phase β into account, see (166) in appendix A.1 and compare to the convention used by the Particle Data Group [71]. In this case the only source of low energy CP violation (and thus in our scenario also CP violation at high energies) is the phase δ. We find

$$\begin{aligned}
\epsilon_1 \;\approx\; & \frac{\tilde{\kappa}^2}{\pi}\,\sin\delta\,\sin\theta_{13}\,\left((-1)^{k_\alpha}\,\sin 2\theta_{23}\,\left[\cos\delta\,\sin\theta_{13}\,\cos 2\theta_{12}\,\sin 2\theta_{23}\right.\right. \qquad (183)\\
& \left.+\frac{1}{2}\,(1+\sin^2\theta_{13})\,\sin 2\theta_{12}\,\cos 2\theta_{23}\right]\,f\left(\frac{m_1}{m_2}\right) + (-1)^{k_\beta}\,\cos\theta_{12}\,\cos^2\theta_{13}\,\cos 2\theta_{23}\,\times\\
& \left.\left[\cos\delta\,\sin\theta_{13}\,\cos\theta_{12}\,\cos 2\theta_{23} - \sin\theta_{12}\,\sin 2\theta_{23}\right]\,f\left(\frac{m_1}{m_3}\right)\right)\\
\epsilon_2 \;\approx\; & \frac{\tilde{\kappa}^2}{\pi}\,(-1)^{k_\alpha}\,\sin\delta\,\sin\theta_{13}\,\left(-\sin 2\theta_{23}\,\left[\cos\delta\,\sin\theta_{13}\,\cos 2\theta_{12}\,\sin 2\theta_{23}\right.\right. \qquad (184)\\
& \left.+\frac{1}{2}\,(1+\sin^2\theta_{13})\,\sin 2\theta_{12}\,\cos 2\theta_{23}\right]\,f\left(\frac{m_2}{m_1}\right) + (-1)^{k_\beta}\,\sin\theta_{12}\,\cos^2\theta_{13}\,\cos 2\theta_{23}\,\times\\
& \left.\left[\cos\delta\,\sin\theta_{13}\,\sin\theta_{12}\,\cos 2\theta_{23} + \cos\theta_{12}\,\sin 2\theta_{23}\right]\,f\left(\frac{m_2}{m_3}\right)\right)\\
\epsilon_3 \;\approx\; & \frac{\tilde{\kappa}^2}{\pi}\,(-1)^{k_\beta}\,\sin\delta\,\sin\theta_{13}\,\cos^2\theta_{13}\,\cos 2\theta_{23}\,\left(\cos\theta_{12}\,\left[\sin\theta_{12}\,\sin 2\theta_{23}\right.\right. \qquad (185)\\
& \left.-\cos\delta\,\sin\theta_{13}\,\cos\theta_{12}\,\cos 2\theta_{23}\right]\,f\left(\frac{m_3}{m_1}\right) + (-1)^{k_\alpha+1}\,\sin\theta_{12}\,\left[\cos\delta\,\sin\theta_{13}\,\sin\theta_{12}\,\cos 2\theta_{23}\right.\\
& \left.\left.+\cos\theta_{12}\,\sin 2\theta_{23}\right]\,f\left(\frac{m_3}{m_2}\right)\right)
\end{aligned}$$

Note that none of the CP asymmetries ϵ_i can be written in a form similar to the one presented in (81). Thus, the sign of ϵ_i cannot be predicted with the knowledge of the sign of $\sin\delta$ (and of the loop function) only. For example, the sign of ϵ_3 also depends on the octant of θ_{23}, $\theta_{23} \lessgtr \pi/4$, since ϵ_3 is proportional to $\cos 2\theta_{23}$. These formulae are consistent with those derived for case 3 b.1), $m = n/2$ and $s = n/2$, see (99-101), since both lead to vanishing CP asymmetries, if we take into account maximal atmospheric mixing and maximal Dirac phase for (183-185) and $\zeta = \pi/2, 3\,\pi/2$ for (99-101).

We also analyze the case in which the Majorana phase α and the Dirac phase δ are trivial

$$\alpha = k_\alpha\,\pi \;\;,\;\; \delta = k_\delta\,\pi \quad \text{with} \quad k_{\alpha,\delta} = 0, 1\;. \tag{186}$$

Like the case with trivial β instead of α, presented in subsection 3.2, the formulae of the CP asymmetries are compact and, especially, we can express ϵ_3 in terms of the other two ones. For ϵ_1 and ϵ_2 we find

$$\begin{aligned}
\epsilon_1 \;\approx\; & -\frac{\tilde{\kappa}^2}{2\,\pi}\,\sin\beta\,\cos^2\theta_{13}\,\left[\sin\theta_{12}\,\sin 2\theta_{23} - (-1)^{k_\delta}\,\sin\theta_{13}\,\cos\theta_{12}\,\cos 2\theta_{23}\right]^2\,f\left(\frac{m_1}{m_3}\right)\;, \qquad (187)\\
\epsilon_2 \;\approx\; & \frac{\tilde{\kappa}^2}{2\,\pi}\,(-1)^{k_\alpha+1}\,\sin\beta\,\cos^2\theta_{13}\,\left[\cos\theta_{12}\,\sin 2\theta_{23} + (-1)^{k_\delta}\,\sin\theta_{13}\,\sin\theta_{12}\,\cos 2\theta_{23}\right]^2\,f\left(\frac{m_2}{m_3}\right)
\end{aligned}$$

and ϵ_3 reads

$$\epsilon_3 = -\epsilon_1 \, f\left(\frac{m_3}{m_1}\right)\left(f\left(\frac{m_1}{m_3}\right)\right)^{-1} - \epsilon_2 \, f\left(\frac{m_3}{m_2}\right)\left(f\left(\frac{m_2}{m_3}\right)\right)^{-1} \, . \tag{188}$$

Similarly, to the case in which α is trivial, the CP asymmetries $\epsilon_{1,2}$ can be decomposed into three pieces: the sine of the non-trivial Majorana phase β, the loop function and a piece that can be written as a square. In this way it becomes evident what determines the sign of these CP asymmetries.

D Sketch of models

In the following we present ideas for explicit realizations of our scenario in non-SUSY as well as SUSY contexts. For the sake of concreteness we will stick to the flavor group $\Delta(6\,n^2)$ with $n = 4$ and choose one particular CP transformation X. In particular, we motivate the size κ and the flavor structure of the correction δY_D. We note that models with the flavor group $\Delta(96)$ and the corresponding group $\Delta(48)$ can be found in the literature [24, 76].

D.1 Non-SUSY setup

Here we present a sketch of a non-SUSY model. Apart from the flavor group $\Delta(96)$ and the auxiliary symmetry $Z_3^{(\mathrm{aux})}$ we also assume the presence of an additional Z_{12} group that we use to segregate better the charged lepton and the neutrino sectors. LH leptons l and RH neutrinos N transform as **3** that is a complex faithful representation of $\Delta(96)$, while RH charged leptons α_R are trivial singlets. The fields l and N are neutral under $Z_3^{(\mathrm{aux})}$, whereas α_R carry the charges 1 for $\alpha = e$, ω for $\alpha = \mu$ and ω^2 for $\alpha = \tau$, respectively. The fields l and N also transform in the same way under the additional Z_{12} group, since they have the charge ω_{12}^4, $\omega_{12} = e^{2\,\pi\,i/12}$. RH charged leptons all are assigned the phase ω_{12}^3. As CP symmetry we impose the one generated by $X = a\,b\,c\,d^2\,P_{23}$ in the representation **3**.[16]

Charged lepton masses are produced with the help of three different flavor symmetry breaking fields ϕ_α that transform as

$$\phi_e \sim (\mathbf{3}, 1, \omega_{12}) \quad , \quad \phi_\mu \sim (\mathbf{3}, \omega^2, \omega_{12}) \quad \text{and} \quad \phi_\tau \sim (\mathbf{3}, \omega, \omega_{12}) \tag{189}$$

under $(\Delta(96), Z_3^{(\mathrm{aux})}, Z_{12})$ and are neutral under the gauge group. The relevant terms in the Lagrangian read

$$\frac{y_e}{\Lambda}\,\bar{l}\,H\,\phi_e\,e_R + \frac{y_\mu}{\Lambda}\,\bar{l}\,H\,\phi_\mu\,\mu_R + \frac{y_\tau}{\Lambda}\,\bar{l}\,H\,\phi_\tau\,\tau_R + \mathrm{h.c.} \tag{190}$$

with Λ being the cutoff scale of the theory and y_α real couplings. The VEVs of ϕ_α are taken to be

$$\langle\phi_e\rangle \propto \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \quad , \quad \langle\phi_\mu\rangle \propto \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \quad \text{and} \quad \langle\phi_\tau\rangle \propto \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \tag{191}$$

and thus break G_f to the residual group $Z_3^{(D)}$, as desired. Since the phases of the VEVs are in general undetermined, they break the CP symmetry imposed on the theory. The charged lepton mass matrix, computed from (190) and (191), is diagonal. For y_τ of order one the correct tau lepton mass is obtained for

$$|\langle\phi_\tau\rangle|\,/\Lambda = \varepsilon \approx (0.01 \div 0.1) \, . \tag{192}$$

Assuming all VEVs $\langle\phi_\alpha\rangle$ to be of that order, the correct hierarchy of the charged lepton masses can be achieved with an additional Froggatt-Nielsen symmetry under which RH charged leptons carry different charges.

[16] The explicit form of X in the other representations of $\Delta(96)$ can be calculated using the information about the corresponding automorphism given in (30).

Since RH neutrinos transform like LH leptons under the flavor and CP symmetry, the neutrino Yukawa coupling Y_D arises from a renormalizable operator. Its flavor structure is trivial and the coupling is real. The Majorana mass term of RH neutrinos originates from couplings to two different triplets of $\Delta(96)$, one equivalent to $\mathbf{3}$ and another one to $\mathbf{3'}$ which is a real and unfaithful representation of $\Delta(96)$,

$$\varphi_\nu \sim (\mathbf{3}, 1, \omega_{12}^4) \quad \text{and} \quad \psi_\nu \sim (\mathbf{3'}, 1, \omega_{12}^4) \; . \tag{193}$$

As indicated, these fields are neutral under $Z_3^{(\text{aux})}$, but carry the charge ω_{12}^4 under Z_{12}, so that the Lagrangian contains the following terms

$$f_1 \, \overline{N^c} \, N \, \varphi_\nu + f_2 \, \overline{N^c} \, N \, \psi_\nu + \text{h.c.} \tag{194}$$

with f_1 and f_2 being real couplings. The VEVs $\langle \varphi_\nu \rangle$ and $\langle \psi_\nu \rangle$ are aligned as follows

$$\langle \varphi_\nu \rangle = i \, w \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \quad , \quad \langle \psi_\nu \rangle = \begin{pmatrix} v_1 + i \, v_2 \\ v_1 + i \, v_3 \\ v_1 - i \, (v_2 + v_3) \end{pmatrix} \tag{195}$$

with $v_{1,2,3}$ and w real parameters of order $\varepsilon^2 \, \Lambda$. Thus, RH neutrino masses between $10^{12}\,\text{GeV}$ and $10^{14}\,\text{GeV}$ are achieved for Λ close to the scale of grand unification. Notice that we have chosen the VEVs of φ_ν and ψ_ν to be smaller than those of the fields ϕ_α. In this way the largest correction to the Dirac neutrino mass matrix arises from the fields ϕ_α only, see (198). As one can check explicitly, $\langle \varphi_\nu \rangle$ and $\langle \psi_\nu \rangle$ leave G_ν, generated by $Z = c^2$ and the CP transformation X, invariant. This breaking pattern hence allows us to obtain the PMNS mixing matrix of case 1), see (31), for $n = 4$ and $s = 1$. This is one choice of parameters also employed in our numerical discussion of unflavored leptogenesis in case 1), see subsection 3.3. The free parameter θ depends on the VEV of the field ψ_ν

$$\tan 2\,\theta = -\frac{v_2 + 2\, v_3}{\sqrt{3}\, v_2} \; . \tag{196}$$

Its particular value, necessary for describing correctly the lepton mixing angles, should be explained in a more complete model. The three RH neutrino masses M_i read[17]

$$\begin{aligned} M_1 &= \sqrt{3} \left| f_1 \, \text{sign}(v_2) \sqrt{v_2^2 + v_2 \, v_3 + v_3^2} + \sqrt{3} \, f_2 \, w \right| \; , \\ M_2 &= 3 \, |f_1 \, v_1| \; , \\ M_3 &= \sqrt{3} \left| f_1 \, \text{sign}(v_2) \sqrt{v_2^2 + v_2 \, v_3 + v_3^2} - \sqrt{3} \, f_2 \, w \right| \end{aligned} \tag{197}$$

and thus are functions of both couplings $f_{1,2}$ as well as all parameters of the VEVs of the fields φ_ν and ψ_ν. Using these we can also compute the masses of the light neutrinos. As one can see, we can accommodate in this way both mass orderings as well as a QD mass spectrum.

The discussed operators necessary at LO in our scenario contain either no or one flavor symmetry breaking field. Each mass matrix, m_l, m_D and M_R, receives corrections of relative order ε^2 with respect to the corresponding LO result. Those to m_l arise from insertions of three fields ϕ_α and are of the generic form $\phi_\alpha \phi_\beta \phi_\gamma^\dagger$ with α, β, γ being e, μ or τ. Clearly, these do not change the form of m_l and thus the charged lepton mass matrix is still diagonal. The dominant corrections to m_D instead change the form of the latter and hence constitute the leading form of δY_D. They stem from the terms

$$\sum_{\alpha=e,\mu,\tau} \sum_{r=1,2,6} \frac{y^\nu_{\alpha,r}}{\Lambda^2} \, \bar{l} \, H^c \, \phi_\alpha^\dagger \, \phi_\alpha \, N + \text{h.c.} \; . \tag{198}$$

The index α indicates which field ϕ_α is coupled, while the index r takes into account the different possible contractions via a one-, two- or six-dimensional representation of $\Delta(96)$. The contribution

[17] $\text{sign}(x)$ stands for the sign of the real parameter x.

arising from the contraction to a singlet can be absorbed into the LO term, since it is always real and proportional to the identity matrix in flavor space. The one to a doublet, indeed, does not contribute, since the residual symmetry G_e that is left invariant by the VEVs of the fields in (191) forces it to vanish. Consequently, the correction δY_D is generated via the terms with $y^\nu_{\alpha,6}$ in (198). Matching the form of δY_D given in (27) the two couplings z_1 and z_2 turn out to be

$$z_1 = \frac{\sqrt{3}}{2}\,\left(2\,y^\nu_{e,6} - y^\nu_{\mu,6} - y^\nu_{\tau,6}\right) \quad \text{and} \quad z_2 = \frac{3}{2}\,\left(y^\nu_{\tau,6} - y^\nu_{\mu,6}\right)\,, \tag{199}$$

if we set $\langle \phi^\dagger_\alpha\,\phi_\alpha\rangle$ to $\varepsilon^2\,\Lambda^2$ for $\alpha = e,\,\mu,\,\tau$ for simplicity. The parameter κ is thus of the order

$$\kappa \approx \varepsilon^2 \quad \text{meaning} \quad 10^{-4} \lesssim \kappa \lesssim 10^{-2}\,. \tag{200}$$

We note that in this particular case $z_{1,2}$ in (199) turn out to be real. Subleading corrections to the Dirac neutrino mass matrix arise from two types of terms: terms with two flavor symmetry breaking fields (φ_ν and ψ_ν and the conjugate fields) and terms with four flavor symmetry breaking fields of the type $\phi^\dagger_\alpha\,\phi^\dagger_\beta\,\phi_\gamma\,\phi_\delta$ with $\alpha,\,\beta,\,\gamma,\,\delta = e,\,\mu,\,\tau$. Both lead to terms relatively suppressed by ε^4 with respect to the LO term. The RH neutrino mass matrix, being at LO of order $\varepsilon^2\,\Lambda$, also receives corrections. The most dominant ones are of order $\varepsilon^4\,\Lambda$ and arise from three types of terms: terms with two conjugate fields $\varphi^\dagger_\nu$ and $\psi^\dagger_\nu$, terms with three fields of the form $\phi^\dagger_\alpha\,\phi_\alpha\,\varphi_\nu$ or $\phi^\dagger_\alpha\,\phi_\alpha\,\psi_\nu$ for $\alpha = e,\,\mu,\,\tau$ as well as terms with four fields of the type ϕ_α that are in general different. Clearly, the latter two types of terms break the residual symmetry G_ν, if the VEVs of the flavor symmetry breaking fields are inserted. These together with the correction δY_D that is also of relative order ε^2 with respect to the corresponding LO term will induce small corrections to the LO results for the lepton mixing parameters. However, these are expected to be suppressed by ε^2 and thus at most at the percent level.

D.2 SUSY setup

If we consider instead a SUSY framework, we can also construct a model of this type. Apart from the fact that l and ν^c transform in complex conjugated three-dimensional representations, $l \sim \overline{\mathbf{3}}$ and $\nu^c \sim \mathbf{3}$, see section 5, the three main differences are: *a*) we slightly change the additional symmetry and we use a Z_5 instead of a Z_{12} group. The transformation properties of the fields are

$$l \sim \omega_5^3\;\;,\;\;\alpha^c \sim \omega_5^2\;\;,\;\;\nu^c \sim \omega_5^2\;\;,\;\;\varphi_\nu\,,\,\psi_\nu \sim \omega_5\quad(\,\omega_5 = e^{2\,\pi\,i/5}\,) \tag{201}$$

and the Higgs supermultiplets h_u and h_d are neutral; *b*) we use less fields in the charged lepton sector

$$\phi_\tau \sim (\mathbf{3},\omega,1) \quad \text{and} \quad \chi \sim (\mathbf{2},\omega,1) \tag{202}$$

under $(\Delta(96), Z_3^{(\text{aux})}, Z_5)$. The VEVs of these fields leave $Z_3^{(D)}$ invariant, if they are chosen to be of the form

$$\langle\phi_\tau\rangle \propto \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \quad \text{and} \quad \langle\chi\rangle \propto \begin{pmatrix} 0 \\ 1 \end{pmatrix}\,. \tag{203}$$

The terms in the superpotential contributing at lowest orders to the charged lepton mass matrix are

$$\frac{y_e}{\Lambda^3}\,l\,h_d\,\phi_\tau\,\chi^2\,e^c + \frac{y_\mu}{\Lambda^2}\,l\,h_d\,\phi_\tau\,\chi\,\mu^c + \frac{y_\tau}{\Lambda}\,l\,h_d\,\phi_\tau\,\tau^c\,. \tag{204}$$

In this way, we can generate the mass hierarchy among the charged leptons with the help of insertions of several flavor symmetry breaking fields. Furthermore, the correct ratio between muon and tau lepton masses and electron and tau lepton masses is achieved, if we assume that the field χ acquires a VEV of the order

$$|\langle\chi\rangle|\,/\Lambda \approx \lambda^2 \approx 0.04\,, \tag{205}$$

while the VEV of ϕ_τ is chosen like in (192) and its actual value depends on the size of $\tan\beta$, see details in section 5. We note also that the VEVs of φ_ν and ψ_ν still have the form as in (195), but we now choose their size to be $\varepsilon\,\Lambda$; *c*) the lowest order correction to the Dirac mass matrix of the neutrinos and thus the source of δY_D are two operators with three insertions of the field ϕ_τ[18]

$$\sum_{i=1,2} \frac{y^\nu_{\tau,i}}{\Lambda^3}\, l\, h_u\, \phi^3_\tau\, \nu^c \tag{206}$$

with $y^\nu_{\tau i}$ being both real. Thus, the size of the small parameter κ is estimated as

$$\kappa \approx \varepsilon^3 \quad \text{meaning} \quad 10^{-6} \lesssim \kappa \lesssim 10^{-3}\ . \tag{207}$$

This has to be compared with the value given in (79) showing that the expansion parameter ε should be in the SUSY realization around $\varepsilon \approx 0.1$. The parameters z_1 and z_2 of δY_D in (27) can be written as

$$z_1 = -\sqrt{3}\,\left(y^\nu_{\tau,1} + y^\nu_{\tau,2}\right) \quad , \quad z_2 = 3\,\left(y^\nu_{\tau,1} + y^\nu_{\tau,2}\right) \tag{208}$$

for $\langle\phi_\tau\rangle$ fixed to $\varepsilon\,\Lambda$.

D.3 General source of correction δY_D

Lastly, we can also consider the general case in which a six-dimensional irreducible representation Φ uncharged under all auxiliary symmetries couples to LH leptons, RH neutrinos and the Higgs field H in a non-SUSY context

$$\frac{y^\nu_\Phi}{\Lambda}\, \bar{l}\, H^c\, \Phi\, N + \text{h.c.}\ . \tag{209}$$

In a SUSY context the corresponding term in the superpotential reads

$$\frac{y^\nu_\Phi}{\Lambda}\, l\, h_u\, \Phi\, \nu^c\ . \tag{210}$$

In both cases y^ν_Φ is a real coupling. The most general form of the VEV of Φ that leaves the residual group $Z_3^{(D)}$ invariant is

$$\langle\Phi\rangle = \begin{pmatrix} x_1 \\ 0 \\ 0 \\ x_2 \\ 0 \\ 0 \end{pmatrix} \kappa\,\Lambda \quad \text{with} \quad x_i \quad \text{complex} \tag{211}$$

and assuming κ like requested in (79). Thus, z_1 and z_2, parametrizing the correction δY_D in (27), read

$$z_1 = \frac{\sqrt{3}}{2}\, y^\nu_\Phi\, (x_1 + x_2) \quad \text{and} \quad z_2 = \frac{\sqrt{3}}{2}\, i\, y^\nu_\Phi\, (x_1 - x_2)\ . \tag{212}$$

This shows that the special cases, $z_1 = 0$ or $z_2 = 0$, discussed in section 3, can be achieved with a particular form of the VEV of the six-dimensional representation Φ. Clearly, the latter can also arise from some combination of flavor symmetry breaking fields.

References

[1] P. A. R. Ade *et al.* [Planck Collaboration], arXiv:1502.01589 [astro-ph.CO].

[2] A. D. Sakharov, Pisma Zh. Eksp. Teor. Fiz. **5** (1967) 32 [JETP Lett. **5** (1967) 24] [Sov. Phys. Usp. **34** (1991) 392] [Usp. Fiz. Nauk **161** (1991) 61].

[18] These two operators correspond to two independent contractions of the flavor indices. Furthermore, we note that the operator with three fields χ gives a vanishing contribution to δY_D, if the VEV of χ, see (203), is inserted.

[3] M. Fukugita and T. Yanagida, Phys. Lett. B **174** (1986) 45.

[4] V. A. Kuzmin, V. A. Rubakov and M. E. Shaposhnikov, Phys. Lett. B **155** (1985) 36.

[5] S. Pascoli, S. T. Petcov and W. Rodejohann, Phys. Rev. D **68** (2003) 093007 [hep-ph/0302054]; J. A. Casas and A. Ibarra, Nucl. Phys. B **618** (2001) 171 [hep-ph/0103065].

[6] T. Yanagida, in *Proceedings of the Workshop on the Unified Theory and the Baryon Number in the Universe* (O. Sawada and A. Sugamoto, eds.), KEK Tsukuba, Japan, 1979, p. 95; S. L. Glashow, *The future of elementary particle physics*, in *Proceedings of the 1979 Cargèse Summer Institute on Quarks and Leptons* (M. Lévy, J.-L. Basdevant, D. Speiser, J. Weyers, R. Gastmans, and M. Jacob, eds.), Plenum Press, New York, 1980, pp. 687-713; M. Gell-Mann, P. Ramond, and R. Slansky, *Complex spinors and unified theories*, in *Supergravity* (P. van Nieuwenhuizen and D. Z. Freedman, eds.), North Holland, Amsterdam, 1979, p. 315; R. N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44** (1980) 912.

[7] K. Abe *et al.* [T2K Collaboration], Phys. Rev. D **91** (2015) 7, 072010 [arXiv:1502.01550 [hep-ex]]; M. R. Salzgeber [T2K Collaboration], arXiv:1508.06153 [hep-ex].

[8] P. Adamson *et al.* [MINOS Collaboration], Phys. Rev. Lett. **112** (2014) 191801 [arXiv:1403.0867 [hep-ex]].

[9] P. Adamson *et al.* [NOvA Collaboration], arXiv:1601.05037 [hep-ex]; P. Adamson *et al.* [NOvA Collaboration], arXiv:1601.05022 [hep-ex].

[10] R. Acciarri *et al.* [DUNE Collaboration], arXiv:1512.06148 [physics.ins-det].

[11] K. Abe *et al.* [Hyper-Kamiokande Working Group Collaboration], arXiv:1412.4673 [physics.ins-det].

[12] W. Rodejohann, Int. J. Mod. Phys. E **20** (2011) 1833 [arXiv:1106.1334 [hep-ph]].

[13] S. Pascoli, S. T. Petcov and A. Riotto, Phys. Rev. D **75** (2007) 083511 [hep-ph/0609125]; S. Pascoli, S. T. Petcov and A. Riotto, Nucl. Phys. B **774** (2007) 1 [hep-ph/0611338]; E. Molinaro, S. T. Petcov, T. Shindou and Y. Takanishi, Nucl. Phys. B **797** (2008) 93 [arXiv:0709.0413 [hep-ph]]; E. Molinaro and S. T. Petcov, Eur. Phys. J. C **61** (2009) 93 [arXiv:0803.4120 [hep-ph]]; E. Molinaro and S. T. Petcov, Phys. Lett. B **671** (2009) 60 [arXiv:0808.3534 [hep-ph]].

[14] M. C. Gonzalez-Garcia, M. Maltoni and T. Schwetz, JHEP **1411** (2014) 052 [arXiv:1409.5439 [hep-ph]].

[15] F. Capozzi, E. Lisi, A. Marrone, D. Montanino and A. Palazzo, arXiv:1601.07777 [hep-ph].

[16] D. V. Forero, M. Tortola and J. W. F. Valle, Phys. Rev. D **90** (2014) 9, 093006 [arXiv:1405.7540 [hep-ph]].

[17] G. Altarelli, F. Feruglio, Rev. Mod. Phys. **82** (2010) 2701-2729 [arXiv:1002.0211 [hep-ph]]; H. Ishimori, T. Kobayashi, H. Ohki, Y. Shimizu, H. Okada, M. Tanimoto, Prog. Theor. Phys. Suppl. **183** (2010) 1-163 [arXiv:1003.3552 [hep-th]]; S. F. King and C. Luhn, Rept. Prog. Phys. **76** (2013) 056201 [arXiv:1301.1340 [hep-ph]].

[18] W. Grimus and P. O. Ludl, J. Phys. A **45** (2012) 233001 [arXiv:1110.6376 [hep-ph]].

[19] W. Grimus and L. Lavoura, JHEP **0508**, 013 (2005) [arXiv:hep-ph/0504153]; W. Grimus and L. Lavoura, Phys. Lett. B **572**, 189 (2003) [arXiv:hep-ph/0305046]; G. Altarelli and F. Feruglio, Nucl. Phys. B **741** (2006) 215 [hep-ph/0512103]; I. de Medeiros Varzielas, S. F. King and G. G. Ross, Phys. Lett. B **644** (2007) 153 [hep-ph/0512313]; X. -G. He, Y. -Y. Keum and R. R. Volkas, JHEP **0604** (2006) 039 [hep-ph/0601001]; E. Ma, Phys. Rev. D **70** (2004)

031901 [hep-ph/0404199]; C. S. Lam, Phys. Lett. B **656** (2007) 193 [arXiv:0708.3665 [hep-ph]]; C. S. Lam, Phys. Rev. Lett. **101** (2008) 121602 [arXiv:0804.2622 [hep-ph]]; C. S. Lam, Phys. Rev. D **78** (2008) 073015 [arXiv:0809.1185 [hep-ph]]; A. Blum, C. Hagedorn and M. Lindner, Phys. Rev. D **77** (2008) 076004 [arXiv:0709.3450 [hep-ph]]; R. de Adelhart Toorop, F. Feruglio and C. Hagedorn, Phys. Lett. B **703** (2011) 447 [arXiv:1107.3486 [hep-ph]]; R. de Adelhart Toorop, F. Feruglio and C. Hagedorn, Nucl. Phys. B **858** (2012) 437 [arXiv:1112.1340 [hep-ph]].

[20] S. F. King, T. Neder and A. J. Stuart, Phys. Lett. B **726** (2013) 312 [arXiv:1305.3200 [hep-ph]]; A. S. Joshipura and K. M. Patel, Phys. Lett. B **727** (2013) 480 [arXiv:1306.1890 [hep-ph]]; M. Holthausen, K. S. Lim and M. Lindner, Phys. Lett. B **721** (2013) 61 [arXiv:1212.2411 [hep-ph]]; C. S. Lam, Phys. Rev. D **87** (2013) 013001 [arXiv:1208.5527 [hep-ph]]; C. Hagedorn, A. Meroni and L. Vitale, J. Phys. A **47** (2014) 055201 [arXiv:1307.5308 [hep-ph]]; R. M. Fonseca and W. Grimus, JHEP **1409** (2014) 033 [arXiv:1405.3678 [hep-ph]].

[21] F. Feruglio, C. Hagedorn and R. Ziegler, JHEP **1307** (2013) 027 [arXiv:1211.5560 [hep-ph]].

[22] P. F. Harrison and W. G. Scott, Phys. Lett. B **535** (2002) 163 [hep-ph/0203209]; P. F. Harrison and W. G. Scott, Phys. Lett. B **547** (2002) 219 [hep-ph/0210197]; W. Grimus and L. Lavoura, Phys. Lett. B **579** (2004) 113 [hep-ph/0305309]; P. F. Harrison and W. G. Scott, Phys. Lett. B **594** (2004) 324 [hep-ph/0403278]; Y. Farzan and A. Y. Smirnov, JHEP **0701** (2007) 059 [hep-ph/0610337].

[23] G. J. Ding, S. F. King, C. Luhn and A. J. Stuart, JHEP **1305** (2013) 084 [arXiv:1303.6180 [hep-ph]]; F. Feruglio, C. Hagedorn and R. Ziegler, Eur. Phys. J. C **74** (2014) 2753 [arXiv:1303.7178 [hep-ph]]; G. J. Ding, S. F. King and A. J. Stuart, JHEP **1312** (2013) 006 [arXiv:1307.4212 [hep-ph]]; C. C. Li and G. J. Ding, Nucl. Phys. B **881** (2014) 206 [arXiv:1312.4401 [hep-ph]]; C. C. Li and G. J. Ding, JHEP **1508** (2015) 017 [arXiv:1408.0785 [hep-ph]].

[24] G. J. Ding and Y. L. Zhou, Chin. Phys. C **39** (2015) 2, 021001 [arXiv:1312.5222 [hep-ph]]; G. J. Ding and Y. L. Zhou, JHEP **1406** (2014) 023 [arXiv:1404.0592 [hep-ph]].

[25] G. J. Ding and S. F. King, Phys. Rev. D **89** (2014) 9, 093020 [arXiv:1403.5846 [hep-ph]].

[26] G. J. Ding, S. F. King and T. Neder, JHEP **1412** (2014) 007 [arXiv:1409.8005 [hep-ph]]; G. J. Ding and S. F. King, Phys. Rev. D **93** (2016) 025013 [arXiv:1510.03188 [hep-ph]].

[27] C. Hagedorn, A. Meroni and E. Molinaro, Nucl. Phys. B **891** (2015) 499 [arXiv:1408.7118 [hep-ph]].

[28] C. C. Li and G. J. Ding, JHEP **1505** (2015) 100 [arXiv:1503.03711 [hep-ph]]; A. Di Iura, C. Hagedorn and D. Meloni, JHEP **1508** (2015) 037 [arXiv:1503.04140 [hep-ph]]; P. Ballett, S. Pascoli and J. Turner, Phys. Rev. D **92** (2015) 9, 093008 [arXiv:1503.07543 [hep-ph]].

[29] G. Ecker, W. Grimus and H. Neufeld, Nucl. Phys. B **247** (1984) 70; G. Ecker, W. Grimus and H. Neufeld, J. Phys. A **20** (1987) L807; H. Neufeld, W. Grimus and G. Ecker, Int. J. Mod. Phys. A **3** (1988) 603.

[30] W. Grimus and M. N. Rebelo, Phys. Rept. **281** (1997) 239 [hep-ph/9506272].

[31] M. Holthausen, M. Lindner and M. A. Schmidt, JHEP **1304** (2013) 122 [arXiv:1211.6953 [hep-ph]]; M. -C. Chen, M. Fallbacher, K. T. Mahanthappa, M. Ratz and A. Trautner, Nucl. Phys. B **883** (2014) 267 [arXiv:1402.0507 [hep-ph]].

[32] E. E. Jenkins and A. V. Manohar, Phys. Lett. B **668** (2008) 210 [arXiv:0807.4176 [hep-ph]].

[33] E. Bertuzzo, P. Di Bari, F. Feruglio and E. Nardi, JHEP **0911** (2009) 036 [arXiv:0908.0161 [hep-ph]].

[34] C. Hagedorn, E. Molinaro and S. T. Petcov, JHEP **0909** (2009) 115 [arXiv:0908.0240 [hep-ph]].

[35] D. Aristizabal Sierra, F. Bazzocchi, I. de Medeiros Varzielas, L. Merlo and S. Morisi, Nucl. Phys. B **827** (2010) 34 [arXiv:0908.0907 [hep-ph]].

[36] G. Altarelli and D. Meloni, J. Phys. G **36** (2009) 085005 [arXiv:0905.0620 [hep-ph]].

[37] F. Bazzocchi, L. Merlo and S. Morisi, Phys. Rev. D **80** (2009) 053003 [arXiv:0902.2849 [hep-ph]].

[38] B. Adhikary and A. Ghosal, Phys. Rev. D **78** (2008) 073007 [arXiv:0803.3582 [hep-ph]].

[39] S. F. King and T. Neder, Phys. Lett. B **736** (2014) 308 [arXiv:1403.1758 [hep-ph]].

[40] C. D. Froggatt and H. B. Nielsen, Nucl. Phys. B **147**, 277 (1979).

[41] G. Altarelli and F. Feruglio, Nucl. Phys. B **741** (2006) 215 [hep-ph/0512103].

[42] Y. Lin, Nucl. Phys. B **813** (2009) 91 [arXiv:0804.2867 [hep-ph]].

[43] S. Pascoli, S. T. Petcov and A. Riotto, Phys. Rev. D **75** (2007) 083511 [hep-ph/0609125]; S. Pascoli, S. T. Petcov and A. Riotto, Nucl. Phys. B **774** (2007) 1 [hep-ph/0611338]; S. Blanchet, P. Di Bari and G. G. Raffelt, JCAP **0703** (2007) 012 [hep-ph/0611337].

[44] G. Engelhard, Y. Grossman, E. Nardi and Y. Nir, Phys. Rev. Lett. **99** (2007) 081802 [hep-ph/0612187]; G. Engelhard, Y. Grossman and Y. Nir, JHEP **0707** (2007) 029 [hep-ph/0702151 [hep-ph]].

[45] L. Covi, E. Roulet and F. Vissani, Phys. Lett. B **384** (1996) 169 [hep-ph/9605319].

[46] G. F. Giudice, A. Notari, M. Raidal, A. Riotto and A. Strumia, Nucl. Phys. B **685** (2004) 89 [hep-ph/0310123].

[47] W. Buchmuller and M. Plumacher, Phys. Lett. B **431** (1998) 354 [hep-ph/9710460].

[48] S. Blanchet and P. Di Bari, JCAP **0606** (2006) 023 [hep-ph/0603107].

[49] S. Davidson, E. Nardi and Y. Nir, Phys. Rept. **466** (2008) 105 [arXiv:0802.2962 [hep-ph]].

[50] A. Abada, S. Davidson, A. Ibarra, F.-X. Josse-Michaux, M. Losada and A. Riotto, JHEP **0609** (2006) 010 [hep-ph/0605281].

[51] A. Abada, S. Davidson, F. X. Josse-Michaux, M. Losada and A. Riotto, JCAP **0604** (2006) 004 [hep-ph/0601083].

[52] E. Nardi, Y. Nir, E. Roulet and J. Racker, JHEP **0601** (2006) 164 [hep-ph/0601084].

[53] S. Davidson and A. Ibarra, Phys. Lett. B **535** (2002) 25 [hep-ph/0202239].

[54] R. N. Mohapatra and C. C. Nishi, JHEP **1508** (2015) 092 [arXiv:1506.06788 [hep-ph]].

[55] S. P. Martin and M. T. Vaughn, Phys. Rev. D **50** (1994) 2282 [Erratum-ibid. D **78** (2008) 039903] [hep-ph/9311340]; J. Hisano, T. Moroi, K. Tobe and M. Yamaguchi, Phys. Rev. D **53** (1996) 2442 [hep-ph/9510309].

[56] F. Borzumati and A. Masiero, Phys. Rev. Lett. **57** (1986) 961.

[57] J. Adam *et al.* [MEG Collaboration], Phys. Rev. Lett. **110** (2013) 201801 [arXiv:1303.0754 [hep-ex]].

[58] P. Wintz, *Prepared for 29th International Conference on High-Energy Physics (ICHEP 98), Vancouver, Canada, 23-29 Jul 1998*

[59] M. Raidal, A. van der Schaaf, I. Bigi, M. L. Mangano, Y. K. Semertzidis, S. Abel, S. Albino and S. Antusch *et al.*, Eur. Phys. J. C **57** (2008) 13 [arXiv:0801.1826 [hep-ph]].

[60] S. Weinberg, Phys. Rev. Lett. **48** (1982) 1303; M. Y. Khlopov and A. D. Linde, Phys. Lett. B **138** (1984) 265; J. R. Ellis, J. E. Kim and D. V. Nanopoulos, Phys. Lett. B **145** (1984) 181.

[61] H. Baer, S. Kraml, A. Lessa and S. Sekmen, JCAP **1104** (2011) 039 [arXiv:1012.3769 [hep-ph]].

[62] W. Rodejohann, Int. J. Mod. Phys. E **20** (2011) 1833 [arXiv:1106.1334 [hep-ph]].

[63] M. Agostini *et al.* [GERDA Collaboration], Phys. Rev. Lett. **111** (2013) 12, 122503 [arXiv:1307.4720 [nucl-ex]].

[64] A. Gando *et al.* [KamLAND-Zen Collaboration], Phys. Rev. Lett. **110** (2013) 6, 062502 [arXiv:1211.3863 [hep-ex]].

[65] J. B. Albert *et al.* [EXO-200 Collaboration], Nature **510** (2014) 229 [arXiv:1402.6956 [nucl-ex]].

[66] K. Alfonso *et al.* [CUORE Collaboration], Phys. Rev. Lett. **115** (2015) 10, 102502 [arXiv:1504.02454 [nucl-ex]].

[67] R. Arnold *et al.* [NEMO 3 Collaboration], Phys. Rev. D **89** (2014) 11, 111101 [arXiv:1311.5695 [hep-ex]].

[68] D. R. Artusa *et al.* [CUORE Collaboration], Adv. High Energy Phys. **2015** (2015) 879871 [arXiv:1402.6072 [physics.ins-det]].

[69] R. Arnold *et al.* [SuperNEMO Collaboration], Eur. Phys. J. C **70** (2010) 927 [arXiv:1005.1241 [hep-ex]].

[70] N. Abgrall *et al.* [Majorana Collaboration], Adv. High Energy Phys. **2014** (2014) 365432 [arXiv:1308.1633 [physics.ins-det]].

[71] K. A. Olive *et al.* [Particle Data Group Collaboration], Chin. Phys. C **38** (2014) 090001.

[72] C. Jarlskog, Phys. Rev. Lett. **55** (1985) 1039.

[73] E. E. Jenkins and A. V. Manohar, Nucl. Phys. B **792** (2008) 187 [arXiv:0706.4313 [hep-ph]].

[74] C. Luhn, S. Nasri and P. Ramond, J. Math. Phys. **48** (2007) 073501 [hep-th/0701188].

[75] J. A. Escobar and C. Luhn, J. Math. Phys. **50** (2009) 013524 [arXiv:0809.0639 [hep-th]].

[76] S. F. King, C. Luhn and A. J. Stuart, Nucl. Phys. B **867** (2013) 203 [arXiv:1207.5741 [hep-ph]].